



SunLong Biotech Co.,LTD
Tel: 0086-571- 56623320 Fax:0086-571- 56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[KD-Validated] Anti-SMYD3 Rabbit Monoclonal Antibody

Cat No.: KD-10823

Aliases:

SMYD3; SET And MYND Domain Containing 3; ZNFN3A1; ZMYND1; KMT3E; Zinc Finger MYND Domain-Containing Protein 1; Histone-Lysine N-Methyltransferase SMYD3 ; SET And MYND Domain-Containing Protein 3; Zinc Finger, MYND Domain Containing 1; Zinc Finger Protein, Subfamily 3A (MYND Domain Containing), 1; EC 2.1.1.354; EC 2.1.1.43; BA74P14.1; EC 2.1.1

Background:

UniProt Entry: [Q9H7B4](#);NCBI Gene Entry: [64754](#)

Application Information

Molecular Weight: Predicted, 49 kDa, observed, 42 kDa Clonality: Rabbit monoclonal antibody Clone ID: 24GB2260 Species Reactivity: Human, Mouse, Rat Applications Tested: Western Blotting (WB), Flow Cytometry (FCM), Immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human SMYD3

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage


Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:1,000-1:5,000 Flow Cytometry (FCM): 1:2,000
Immunocytochemistry (IC): 1:1,000

Protocols


For general and specific antibody protocols please visit our website. Read all instructions before using this product.




Flow cytometric analysis of SMYD3 expression in HepG2 cells using anti-SMYD3 antibody 1:2,000. Green, isotype control; red, SMYD3.



Immunocytochemical staining of HepG2 cells with anti-SMYD3 antibody 1:1,000. Nuclei were stained blue with DAPI; SMYD3 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μ m.



Western blotting analysis using anti-SMYD3 antibody 1:5,000 and HRP-conjugated goat anti-rabbit secondary antibody 1:20,000 respectively. Image was developed using FeQ™ ECL Substrate Kit .



Western blotting analysis using anti-SMYD3 antibody 1:5,000 and HRP-conjugated goat anti-rabbit secondary antibody 1:20,000 respectively. Image was developed using NaQ™ ECL Substrate Kit .