

Goat Anti-Mouse IgG H&L / APC antibody

SL0296G-APC

Product Name Goat Anti-Mouse IgG H&L / APC
Chinese Name APC 标记的羊抗小鼠 IgG H&L
Alias Goat Anti-Mouse IgG H&L (APC); Immunoglobulin G;

Specific References (1) | SL0296G-APC has been referenced in 1 publications.



[IF=3.575] Huayuan Liu. et al. FNDC5 induces M2 macrophage polarization and promotes hepatocellular carcinoma cell growth by affecting the PPAR γ /NF- κ B/NLRP3 pathway. Biochem Bioph Res Co. 2021 Dec;582:77
IF ; Human.

PubMed:34695754

Immunogen Species Goat
Clonality Polyclonal
React Species Mouse,
 Flow-Cyt=1:100-1000 IF=1:100-1000
Applications IF=1:100-1000,Flow-Cyt=1:100-1000,ICC/IF=1:100-1000 not yet tested in other applications.
 optimal dilutions/concentrations should be determined by the end user.
Form Liquid
Concentration 2.0 mg/ml
immunogen Native Mouse IgG
Lsotype IgG
Purification affinity purified by Protein G, nonspecific adsorbed
Buffer Solution 10 mM TBS (pH=7.4) with 1% BSA, 3% Proclin300 and 50% glycerol.
Storage Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Attention This product as supplied is intended for research use only, not for use in



Product Detail

human, therapeutic or diagnostic applications.

Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10¹¹ variants.