

Rabbit Anti-Mouse IgG H&L / Cy3 antibody

SL0296R-Cy3

Product Name Rabbit Anti-Mouse IgG H&L / Cy3
Chinese Name Cy3 标记的兔抗小鼠 IgG H&L
Alias Rabbit Anti-Mouse IgG H&L (Cy3); Immunoglobulin G;

Specific References (3) | SL0296R-Cy3 has been referenced in 3 publications.

[IF=6.78] Ning Zhang. et al. Spatial transcriptomics shows moxibustion promotes hippocampus astrocyte and neuron interaction. LIFE SCI. 2022 Oct;;121052 **IF** ; **Mouse**.

PubMed:36220370



[IF=6.208] Ting Lin. et al. CTH/H2S Regulates LPS-Induced Inflammation through IL-8 Signaling in MAC-T Cells. INT J MOL SCI. 2022 Jan;23(19):11822 **IF** ; **Mouse**.

PubMed:36233122

[IF=1.076] Lv et al. Production and immunogenicity of chimeric virus-like particles containing the spike glycoprotein of infectious bronchitis virus. (2014) J.Vet.Sc. 15:209-16 **IF(ICC)** ; **Mouse**.

PubMed:24378590

Immunogen Species Rabbit
Clonality Polyclonal
React Species Mouse,
 IF=1:100-1000,ICC/IF=1:100-1000,Flow-Cyt=1:100-1000
Applications not yet tested in other applications.
 optimal dilutions/concentrations should be determined by the end user.
Form Liquid



| | |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Concentration | 2.0 mg/ml |
| immunogen | Native Mouse IgG |
| Lsotype | IgG |
| Purification | affinity purified by Protein A |
| Buffer Solution | 10 mM TBS (pH=7.4) with 1% BSA, 3% Proclin300 and 50% glycerol. |
| Storage | Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Attention | <p>This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</p> <p>Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10¹¹ variants.</p> |
| Product Detail | |