

## Goat Anti-Human IgG H&L / Cy3 antibody

SL0297G-Cy3

**Product Name** Goat Anti-Human IgG H&L / Cy3  
**Chinese Name** Cy3 标记的羊抗人 IgG H&L  
**Alias** Goat Anti-Human IgG H&L (Cy3); Immunoglobulin G;

**Specific References (1)** | SL0297G-Cy3 has been referenced in 1 publications.



**[IF=0]** He M et al. A Femtoliter On-Demand Droplet Dispensing System Based on Gigahertz Acoustic Resonator.Lab Chip. 2018 Aug 21;18(17):2540-2546. **Other ; Human.**

PubMed:30043817

**Immunogen Species** Goat  
**Clonality** Polyclonal  
**React Species** Human,  
IF=1:100-1000,ICC/IF=1:100-1000,Flow-Cyt=1:100-1000  
**Applications** not yet tested in other applications.  
optimal dilutions/concentrations should be determined by the end user.  
**Form** Liquid  
**Concentration** 2.0 mg/ml  
**immunogen** Native Human IgG  
**Lsotype** IgG  
**Purification** affinity purified by Protein G, nonspecific adsorbed  
**Buffer Solution** 10 mM TBS (pH=7.4) with 1% BSA, 3% Proclin300 and 50% glycerol.  
**Storage** Store at -20°C for one year. Avoid repeated freeze/thaw cycles.  
**Attention** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.  
**Product Detail** Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our



defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10<sup>11</sup> variants.