

Rabbit Anti-Bovine IgM / RBITC antibody

SL0327R-RBITC

Product Name	Rabbit Anti-Bovine IgM / RBITC
Chinese Name	罗丹明标记的兔抗牛 IgM
Alias	Rabbit Anti-Bovine IgM (RBITC); Immunoglobulin M;
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Cow, IF=1:100-1000
Applications	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Form	Liquid
Concentration	2.0 mg/ml
immunogen	Native Bovine IgG
Lsotype	IgG
Purification	affinity purified by Protein A
Buffer Solution	10 mM TBS (pH=7.4) with 1% BSA, 3% Proclin300 and 50% glycerol.
Storage	Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. IgM normally constitutes about 10% of serum immunoglobulins. IgM antibody is prominent in early immune responses to most antigens and is largely confined to plasma due to it's large size. Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. Due to it's high valency IgM is more efficient than other isotypes is binding antigens with repeating epitopes (virus particles and red blood cells) and is more efficient than IgG in activating the complement pathway. The gene for the mu constant region contains four domains separated by short intervening sequences.
Product Detail	