



Goat Anti-Mouse IgM antibody

SL0368G

Product Name:	Goat Anti-Mouse IgM
Chinese Name:	羊抗小鼠IgM
Alias:	Immunoglobulin M.
文献引用 PubMed :	<p>Specific References(2) SL0368G has been referenced in 2 publications.</p> <p>[IF=3.73]Zhao, Zhiyi, et al. "Maxadilan prevents apoptosis in iPS cells and shows no effects on the pluripotent state or karyotype." PloS one 7.3 (2012): e33953.Human. PubMed:22457805</p> <p>[IF=5.44]Zhang, Xiaoru, et al. "Ultrasensitive Photoelectrochemical Immunoassay of Antibody against Tumor-associated Carbohydrate antigen Amplified by Functionalized Graphene Derivates and Enzymatic Biocatalytic Precipitation." Biosensors and Bioelectronics (2014).other; PubMed:not posted yet</p>
Organism Species:	Goat
Clonality:	Polyclonal
React Species:	mo
Applications:	WB=1:1000-5000ELISA=1:2000-10000IHC-P=1:200-1000IHC-F=1:200-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	900kDa
Form:	Lyophilized or Liquid
Concentration:	10mg/1ml
immunogen:	Full length plasma protein:
Lsotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M PBS(pH7.4)

Storage:	Storage: Store at –20 oC for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20oC. When reconstituted in sterile distilled water or diluent supplied, the antibody is stable for at least two weeks at 2-4 °C.
Product Detail:	<p>Immunoglobulin M (IgM) normally constitutes about 10% of serum immunoglobulins. IgM antibody is prominent in early immune responses to most antigens and is largely confined to plasma due to its large size. Monomeric IgM is expressed as a membrane bound antibody on the surface of B cells and as a pentamer when secreted by plasma cells. Due to its high valency IgM is more efficient than other isotypes in binding antigens with repeating epitopes (virus particles and red blood cells) and is more efficient than IgG in activating the complement pathway. The gene for the mu constant region contains four domains separated by short intervening sequences.</p> <p>Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.</p>

www.sunlongbiotech.com