

## Rabbit Anti-DMBT1 antibody

SL0595R

**Product Name** DMBT1

**Chinese Name** 抑癌基因抗体

**Alias** GP340; Apactin; CRP; CRP-[a]; CRP-[b]; CRP-ductin; Crpd; DBMT1; Deleted in malignant brain tumors 1; Deleted in malignant brain tumors 1 protein; DMBT 1; DMBT1; DMBT1 prototype precursor; DMBT1/6kb.1 protein precursor; DMBT1/8kb.1 protein; DMBT1/8kb.2 protein precursor; Ebnerin; Glycoprotein 300; Glycoprotein 340; Gp 340; gp300; Gp340 variant protein; Hensin; Mucin-like glycoprotein; Muclin; p80; Pancrin; Salivary agglutinin; Surfactant pulmonary associated D binding protein; Surfactant pulmonary associated protein D binding protein; Vomeroglandin.

**Research Area** Tumour Cell biology Cell type markers

**Immunogen Species** Rabbit

**Clonality** Polyclonal

**React Species** Mouse, (predicted: Human, Rat, )

**Applications** IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 (Paraffin sections need antigen repair)  
not yet tested in other applications.  
optimal dilutions/concentrations should be determined by the end user.

**Theoretical molecular weight** 156-265kDa

**Cellular localization** The cell membrane Secretory protein

**Form** Liquid

**Concentration** 1mg/ml

**immunogen** KLH conjugated synthetic peptide derived from human DMBT1: 1701-1785/1785

**Lsotype** IgG

**Purification** affinity purified by Protein A

**Buffer Solution** 1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.



---

<b>Storage</b>	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
<b>Attention</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>PubMed</b>	<a href="#">PubMed</a> DMBT 1 may play roles in mucosal defense system, cellular immune defense and epithelial differentiation. May play a role in liver regeneration. May be an important factor in fate decision and differentiation of transit-amplifying ductular(oval) cells within the hepatic lineage. May function as a binding protein in saliva for the regulation of taste sensation. May play a role as an opsonin receptor for SFTPD and SPAR in macrophages tissues throughout the body, including epithelial cells lining the gastrointestinal tract. Required for terminal differentiation of columnar epithelial cells during early embryogenesis. [Subcellular Location] Secreted. Membrane. [Tissue Specificity] Expressed in von Ebner glands (VEG) (at protein level), olfactory epithelium and the lateral nasal gland. Expressed in transit-amplifying ductular (oval) cells. Up-regulated at day 3 after hepatectomy. Expressed in newly formed bile ducts and in structures resembling intestinal epithelium. Belongs to the DMBT1 family.
<b>Product Detail</b>	<p><b>Function:</b> May be considered as a candidate tumor suppressor gene for brain, lung, esophageal, gastric, and colorectal cancers. May play roles in mucosal defense system, cellular immune defense and epithelial differentiation. May play a role as an opsonin receptor for SFTPD and SPAR in macrophage tissues throughout the body, including epithelial cells lining the gastrointestinal tract. May play a role in liver regeneration. May be an important factor in fate decision and differentiation of transit-amplifying ductular (oval) cells within the hepatic lineage. Required for terminal differentiation of columnar epithelial cells during early embryogenesis. May function as a binding protein in saliva for the regulation of taste sensation. Binds to HIV-1 envelope protein and has been shown to both inhibit and facilitate viral transmission. Displays a broad calcium-dependent binding spectrum against both Gram-positive and Gram-negative bacteria, suggesting a role in defense against bacterial pathogens. Binds to a range of poly-sulfated and poly-phosphorylated ligands which may explain its broad bacterial-binding specificity. Inhibits cytoinvasion of S.enterica. Associates with the actin cytoskeleton and is involved in its remodeling during regulated exocytosis. Interacts with pancreatic zymogens in a pH-dependent manner and may act as a Golgi cargo receptor in the regulated secretory pathway of the pancreatic acinar cell.</p> <p><b>Subunit:</b> Binds SFTPD and SPAR in a calcium-dependent manner. Interacts with LGALS3.</p> <p><b>Subcellular Location:</b> Secreted. Some isoforms may be membrane-bound. Localized to the luminal aspect of crypt cells in the small intestine. In the colon, seen in the luminal aspect of surface epithelial cells. Formed in the ducts of von Ebner gland, and released into the fluid</p>

bathing the taste buds contained in the taste papillae.

**Tissue Specificity:**

Highly expressed in alveolar and macrophage tissues. In some macrophages, expression is seen on the membrane, and in other macrophages, strongly expressed in the phagosome/phagolysosome compartments. Expressed in lung, trachea, salivary gland, small intestine and stomach. In pancreas, expressed in certain cells of the islets of Langerhans. In digestive tract, confined to tissues with large epithelial surfaces. In intestinal tissue, moderately expressed in epithelial cells of the midcrypts and the crypt base. Expression is significantly elevated in intestinal tissue from patients with inflammatory bowel disease (IBD), particularly in surface epithelial and Paneth cells, but not in IBD patients with mutant NOD2. Present in crypt bases of the duodenum, in crypt tops of the colon, and in collecting ducts of the cortical kidney. Expressed in stratified squamous epithelium of vagina and in outer luminal surface and basilar region of columnar epithelial cells in cervix (at protein level). Isoform 1 is secreted to the lumen of the respiratory tract.

**Post-translational modifications:**

Highly N- and O-glycosylated. The O-glycans are heavily sulfated.

**DISEASE:**

Defects in DMBT1 are involved in the development of glioma (GLM) [MIM:137800]. Gliomas are central nervous system neoplasms derived from glial cells and comprise astrocytomas, glioblastoma multiforme, oligodendrogliomas, and ependymomas. Note=Homozygous deletions may be the predominant mechanism of DMBT1 inactivation playing a role in carcinogenesis. DMBT1 is deleted in medulloblastoma and glioblastoma cell lines; point mutations have also been reported in patients with glioma. A loss or reduction of DMBT1 expression has been seen in esophageal, gastric, lung and colorectal carcinomas as well.

**Similarity:**

Belongs to the DMBT1 family.  
Contains 2 CUB domains.  
Contains 14 SRCR domains.  
Contains 1 ZP domain.

**SWISS:**

Q9UGM3

**Gene ID:**

1755

**Database links:**

[Entrez Gene: 1755](#) Human


[Omim: 601969](#) Human

[SwissProt: Q9UGM3](#) Human

[Unigene: 279611](#) Human

DMBT1 是分泌型的清道夫受体，富含半胱氨酸，主要在肠道表达，参与再生和防御病原体的功能。有学者认为：DMBT1 在消化系统的恶性 Tumour 的发生、发展及转移中起一定作用。

**Product  
Picture**



Tissue/cell: mouse stomach tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 1M, pH 6.0 ), Boiling bathing for 15min; Block



endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer

(normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-GP340/DMBT1 Polyclonal Antibody, Unconjugated(SL0595R)

1:200, overnight at 4°C, followed by conjugation to the secondary

antibody(SP-0023) and DAB(C-0010) staining