

Rabbit Anti-KCNMA1/BK channel antibody

SL0729R

Product Name KCNMA1/BK channel

Chinese Name 钙激活钾 Channel protein α 1 抗体

Alias Maxi Potassium channel alpha; bA205K10.1; BK channel; BKCa channels; BKCA alpha; BKCA alpha subunit; BKTM; Calcium activated potassium channel subfamily M subunit alpha 1; Calcium activated potassium channel subunit alpha 1; DKFZp686K1437; Drosophila slowpoke like; hSlo; K(VCA)alpha; KCa1.1; KCNMA 1; KCNMA; KCNMA1; Large conductance calcium activated potassium channel subfamily M alpha member 1; Maxi K; Maxi K channel; MaxiK; Potassium large conductance calcium activated channel subfamily M alpha member 1; SAKCA; Slo 1; SLO alpha; SLO; Slo homolog; Slo1; Slowpoke homolog; Stretch activated Kca channel; KCMA1_HUMAN.

Research Area Neurobiology Signal transduction Channel protein

Immunogen Species Rabbit

Clonality Polyclonal

React Species Rat, (predicted: Human, Mouse, Dog, Pig, Cow, Rabbit, Sheep,)
IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 (Paraffin sections need antigen repair)

Applications not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.

Theoretical molecular weight 137kDa

Cellular localization The cell membrane

Form Liquid

Concentration 1mg/ml

immunogen KLH conjugated synthetic peptide derived from human BK channel: 1131-1236/1236

Lsotype IgG

Purification affinity purified by Protein A

Buffer 1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.

**Solution****Storage**

Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

Attention

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

PubMed

[PubMed](#)

MaxiK channels are large conductance, voltage and calcium-sensitive potassium channels which are fundamental to the control of smooth muscle tone and neuronal excitability. MaxiK channels can be formed by 2 subunits: the pore-forming alpha subunit, which is the product of this gene, and the modulatory beta subunit.

Intracellular calcium regulates the physical association between the alpha and beta subunits. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008].

Product Detail**Function:**

Potassium channel activated by both membrane depolarization or increase in cytosolic Ca(2+) that mediates export of K(+). It is also activated by the concentration of cytosolic Mg(2+). Its activation dampens the excitatory events that elevate the cytosolic Ca(2+) concentration and/or depolarize the cell membrane. It therefore contributes to repolarization of the membrane potential. Plays a key role in controlling excitability in a number of systems, such as regulation of the contraction of smooth muscle, the tuning of hair cells in the cochlea, regulation of transmitter release, and innate immunity. In smooth muscles, its activation by high level of Ca(2+), caused by ryanodine receptors in the sarcoplasmic reticulum, regulates the membrane potential. In cochlea cells, its number and kinetic properties partly determine the characteristic frequency of each hair cell and thereby helps to establish a tonotopic map. Kinetics of KCNMA1 channels are determined by alternative splicing, phosphorylation status and its combination with modulating beta subunits. Highly sensitive to both iberiotoxin (IbTx) and charybdotoxin (CTX).

Subunit:

Homotetramer.

Subcellular Location:

Membrane; Multi-pass membrane protein.

Tissue Specificity:

Widely expressed. Except in myocytes, it is almost ubiquitously expressed.

Post-translational modifications:

Phosphorylated (Probable). Phosphorylation by kinases such as PKA and/or PKG. In smooth muscles, phosphorylation affects its activity.

DISEASE:

Defects in KCNMA1 are the cause of generalized epilepsy and paroxysmal dyskinesia (GEPD) [MIM:609446]. Epilepsy is one of the most common and debilitating neurological disorders. Paroxysmal dyskinesias are neurological disorders characterized by sudden, unpredictable, disabling attacks of involuntary movement often requiring life-long treatment. The coexistence of epilepsy and paroxysmal dyskinesia in the same individual or family is an increasingly recognized phenomenon. Patients manifest absence seizures, generalized tonic-clonic seizures, paroxysmal nonkinesigenic dyskinesia, involuntary dystonic or choreiform movements. Onset is usually in childhood and patients may have seizures only, dyskinesia only, or both.

Similarity:

Belongs to the potassium channel family. Calcium-activated (TC 1.A.1.3) subfamily. KCa1.1/KCNMA1 sub-subfamily.
Contains 1 RCK N-terminal domain.

SWISS:

Q12791

Gene ID:

3778

Database links:

[Entrez Gene: 3778](#) Human

[Entrez Gene: 16531](#) Mouse

[Entrez Gene: 83731](#) Rat

[Omim: 600150](#) Human

[SwissProt: Q12791](#) Human

[SwissProt: Q08460](#) Mouse

[SwissProt: Q62976](#) Rat

[Unigene: 144795](#) Human

[Unigene: 343607](#) Mouse


[Unigene: 486347](#) Mouse

[Unigene: 30616](#) Rat

Channel protein (Channel Protein)

该抗体主要用于 The cell membrane 通道的研究。

**Product
Picture**



Tissue/cell: rat stomach tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (1M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-BK channel/Maxi Potassium channel alpha Polyclonal Antibody, Unconjugated(SL0729R) 1:100, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining