

Rabbit Anti-Cyclooxygenase 2 , Alexa Fluor® 680 conjugated antibody

SL0732R-AF680

Product Name	Cyclooxygenase 2, Bodipy Fluor 680 conjugated
Chinese Name	AF680 标记的环氧合酶 2/前列腺素内过氧化物合成酶 2 抗体
Alias	PTGS2; COX 2; COX2; COX-2; Cyclooxygenase; Cyclooxygenase2; Cyclooxygenase-2; hCox 2; PGG/HS; PGH synthase 2; PGHS 2; PGHS2; PHS 2; PHS II; PHS2 ; Prostaglandin endoperoxide synthase 2; Prostaglandin G/H synthase 2 precursor; Prostaglandin G/H synthase and cyclooxygenase; Prostaglandin G/H synthase 2; Prostaglandin H2 synthase 2; TIS10II; PGH2_HUMAN.
Research Area	Tumour Cardiovascular immunology Signal transduction Synthesis and Degradation The new supersedes the old
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Human,Mouse
Applications	Flow-Cyt=1µg/Test,IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Theoretical molecular weight	65kDa
Cellular localization	cytoplasmic
Form	Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human Cyclooxygenase 2: 501-604/604
Lsotype	IgG
Purification	affinity purified by Protein A
Buffer Solution	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

Attention

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

PubMed

[PubMed](#)

Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq, Feb 2009]

Function:

Mediates the formation of prostaglandins from arachidonate. May have a role as a major mediator of inflammation and/or a role for prostanoid signaling in activity-dependent plasticity.

Subunit:

Homodimer.

Product Detail

Subcellular Location:

Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein.

Post-translational modifications:

S-nitrosylation by NOS2 (iNOS) activates enzyme activity. S-nitrosylation may take place on different Cys residues in addition to Cys-561.

Similarity:

Belongs to the prostaglandin G/H synthase family. Contains 1 EGF-like domain.

SWISS:

P35354

Gene ID:

5743