

Rabbit Anti-intestinal FABP antibody

SL0898R

Product Name	intestinal FABP
Chinese Name	小肠型脂肪酸 Binding protein 抗体
Alias	FABP 2; FABP2; FABPI; Fatty acid binding protein 2 intestinal; Fatty acid binding protein intestinal; Fatty acid-binding protein; I-FABP; I FABP; IFABP; Intestinal fatty acid binding protein 2; MGC133132; FABPI_HUMAN; Fatty acid binding protein intestinal; Fatty acid-binding protein 2; Fatty acid-binding protein; Intestinal fatty acid binding protein 2; Intestinal-type fatty acid-binding protein; OTTHUMP00000163925.
Research Area	immunology Growth factors and hormones Kinases and Phosphatases Diabetes
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	(predicted: Human, Mouse, Rat, Chicken,) IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Applications	(Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Theoretical molecular weight	28kDa
Cellular localization	cytoplasmic The cell membrane
Form	Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human IFABP: 31-132/132
Lsotype	IgG
Purification	affinity purified by Protein A
Buffer Solution	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

PubMed

[PubMed](#)

FABP are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters. FABP2 is probably involved in triglyceride-rich lipoprotein synthesis. Binds saturated long-chain fatty acids with a high affinity, but binds with a lower affinity to unsaturated long-chain fatty acids. FABP2 may also help maintain energy homeostasis by functioning as a lipid sensor (By similarity). [Subcellular Location] Cytoplasm (By similarity). [Similarity] Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.

Function:

FABP are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters. FABP2 is probably involved in triglyceride-rich lipoprotein synthesis. Binds saturated long-chain fatty acids with a high affinity, but binds with a lower affinity to unsaturated long-chain fatty acids. FABP2 may also help maintain energy homeostasis by functioning as a lipid sensor.

Subcellular Location:

Cytoplasm.

Product Detail

Tissue Specificity:

Expressed in the small intestine and at much lower levels in the large intestine. Highest expression levels in the jejunum.

Similarity:

Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.

SWISS:

P12104

Gene ID:

2169

Database links:

[Entrez Gene: 2169](#) Human

[Entrez Gene: 14079](#) Mouse

[Entrez Gene: 25598](#) Rat

[Omim: 134640](#) Human

[SwissProt: P12104](#) Human

[SwissProt: P55050](#) Mouse

[SwissProt: Q53YP5](#) Mouse

[SwissProt: P02693](#) Rat

[Unigene: 282265](#) Human

[Unigene: 28398](#) Mouse

[Unigene: 91358](#) Rat

小肠脂肪酸 Binding protein(IFABP)主要是在小肠分泌的一种蛋白质,参与了脂肪酸的吸收和代谢。IFABP 有可能通过干扰脂肪酸代谢而影响胰岛素敏感性。 IFABP 主要用于 2 型 Diabetes 相关性的研究。