

Rabbit Anti-phospho-PTPN6 (Tyr564)antibody

SL10167R

Product Name phospho-PTPN6 (Tyr564)

Chinese Name 磷酸化蛋白酪氨酸磷酸酶 1C 抗体

Alias

SHP-1(Phospho-Tyr564); PTPN6(phospho Y564); SHP1(phospho Y564); SHP1; 70 kda SHP 1L protein; 70 kda SHP1L protein; 70Z-SHP; EC 3.1.3.48; HCP; HCPH; Hematopoietic cell phosphatase; Hematopoietic cell protein tyrosine phosphatase; Hematopoietic cell protein-tyrosine phosphatase; HPTP 1C; HPTP1C; MGC124580; Protein tyrosine phosphatase 1C; Protein tyrosine phosphatase non receptor type 6; Protein tyrosine phosphatase SHP 1; Protein tyrosine phosphatase SHP1; Protein-tyrosine phosphatase 1C; Protein-tyrosine phosphatase SHP-1; PTN6_HUMAN; PTP 1C; PTP-1C; PTP1C; PTPN 6; PTPN6; SH PTP 1; SH PTP1; SH-PTP1; SHP 1; SHP 1L; SHP1L; SHPTP 1; SHPTP1; Tyrosine protein phosphatase non receptor type 6; Tyrosine-protein phosphatase non-receptor type 6.

Product Type Phosphorylated anti

Immunogen Species Rabbit

Clonality Polyclonal

React Species Rat (predicted:Human,Mouse,Dog,Pig,Horse,Sheep,GuineaPig)

Applications

WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 (Paraffin sections need antigen repair)
not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.

Theoretical molecular weight 65kDa

Cellular localization The nucleus cytoplasmic

Form Liquid

Concentration 1mg/ml

immunogen KLH conjugated Synthesised phosphopeptide derived from human PTPN6 around the phosphorylation site of Tyr564: DV(p-Y)EN

Lsotype IgG



Purification	affinity purified by Protein A
Buffer Solution	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
PubMed	PubMed The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. N-terminal part of this PTP contains two tandem Src homolog (SH2) domains, which act as protein phospho-tyrosine binding domains, and mediate the interaction of this PTP with its substrates. This PTP is expressed primarily in hematopoietic cells, and functions as an important regulator of multiple signaling pathways in hematopoietic cells. This PTP has been shown to interact with, and dephosphorylate a wide spectrum of phospho-proteins involved in hematopoietic cell signaling. Multiple alternatively spliced variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jul 2008]
Product Detail	Function: Modulates signaling by tyrosine phosphorylated cell surface receptors such as KIT and the EGF receptor/EGFR. The SH2 regions may interact with other cellular components to modulate its own phosphatase activity against interacting substrates. Together with MTUS1, induces UBE2V2 expression upon angiotensin II stimulation. Plays a key role in hematopoiesis. Subunit: Monomer. Interacts with MTUS1. Interacts with MILR1 (tyrosine-phosphorylated). Interacts with KIT. Binds PTPNS1, LILRB1 and LILRB2. Interacts with FCRL2, FCRL3, FCRL4, CD300LF, CDK2 and CD84. Interacts with KIR2DL1; the interaction is enhanced by ARRB2. Interacts (via SH2 1 domain) with ROS1; the interaction is direct and promotes ROS1 dephosphorylation. Interacts with EGFR; inhibits EGFR-dependent activation of MAPK/ERK. Interacts with LYN. Interacts with the tyrosine phosphorylated form of PDPK1. Subcellular Location: Cytoplasm. Nucleus. Note=In neurons, translocates into the nucleus after treatment with angiotensin II. Shuttles between the cytoplasm and nucleus via its association with PDPK1. Tissue Specificity: Isoform 1 is expressed in hematopoietic cells. Isoform 2 is expressed in non-hematopoietic cells.

Post-translational modifications:

Phosphorylated on tyrosine residues. Binding of KITLG/SCF to KIT increases tyrosine phosphorylation. Phosphorylation at Tyr-564 enhances phosphatase activity.

Similarity:

Belongs to the protein-tyrosine phosphatase family. Non-receptor class 2 subfamily. Contains 2 SH2 domains.

Contains 1 tyrosine-protein phosphatase domain.

SWISS:

P29350

Gene ID:

5777

Database links:

[Entrez Gene: 5777](#) Human

[Entrez Gene: 15170](#) Mouse

[Entrez Gene: 116689](#) Rat

[Omim: 176883](#) Human

[SwissProt: P29350](#) Human

[SwissProt: P29351](#) Mouse


[SwissProt: P81718](#) Rat

[Unigene: 63489](#) Human

[Unigene: 271799](#) Mouse

[Unigene: 18985](#) Rat

Product
Picture




Sample: Bladder (Rat) Lysate at 40 ug

Primary: Anti-phospho-PTPN6(Tyr564) (SL10167R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 65 kD

Observed band size: 60 kD



Tissue/cell: Rat spleen tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (1M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-phospho-PTPN6 (Tyr564) Polyclonal Antibody,

Unconjugated(SL10167R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining