

Rabbit Anti-Phospho-HDAC5 (Ser259)antibody

SL10326R

Product Name Phospho-HDAC5 (Ser259)

Chinese Name 磷酸化组蛋白去乙酰化酶 5 抗体

Alias HDAC5 (phospho S259); p-HDAC5 (phospho S259); HDAC5(Phospho-Ser259); HDAC5(Phospho-S259); p-HDAC5(Ser259); p-HDAC5(S259); HD 5; HD5; HDAC 5; Histone deacetylase 5; KIAA0600; NY CO 9; Antigen NY CO 9; FLJ90614; HDAC5_HUMAN; Antigen NY-CO-9.

Product Type Phosphorylated anti

Research Area Tumour Cell biology Neurobiology Signal transduction Apoptosis

Immunogen Species Rabbit

Clonality Polyclonal

React Species Human, Mouse, Rat, (predicted: Dog, Pig, Cow, Horse, Rabbit, Sheep,)
WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 (Paraffin sections need antigen repair)
Applications not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.

Theoretical molecular weight 123kDa

Cellular localization The nucleus cytoplasmic

Form Liquid

Concentration 1mg/ml

immunogen KLH conjugated synthesised phosphopeptide derived from human HDAC5 around the phosphorylation site of Ser259 : TA(p-S)EP

Lsotype IgG

Purification affinity purified by Protein A

Buffer Solution 1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.

Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
PubMed	PubMed Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008].
	Function: Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation by repressing transcription of myocyte enhancer MEF2C. During muscle differentiation, it shuttles into the cytoplasm, allowing the expression of myocyte enhancer factors.
Product Detail	Subunit: Interacts with AHRR. Interacts with BAHD1, BCOR, HDAC7, HDAC9, CTBP1, MEF2C, NCOR2, NRIP1, PHB2 and a 14-3-3 chaperone protein. Interacts with KDM5B. Interacts with MYOC. Interacts with GRK5. Interacts with DDIT3/CHOP.
	Subcellular Location: Nucleus. Cytoplasm. Note=Shuttles between the nucleus and the cytoplasm. In muscle cells, it shuttles into the cytoplasm during myocyte differentiation. The export to cytoplasm depends on the interaction with a 14-3-3 chaperone protein and is due to its phosphorylation at Ser-259 and Ser-498 by AMPK, CaMK1 and SIK1.
	Tissue Specificity: Ubiquitous.
	Post-translational modifications: Phosphorylated by AMPK, CaMK1, SIK1 and PRKD1 at Ser-259 and Ser-498. The phosphorylation is required for the export to the cytoplasm and inhibition. Phosphorylated by the PKC kinases PKN1 and PKN2, impairing nuclear import. Phosphorylated by GRK5, leading to nuclear export of HDAC5 and allowing MEF2-mediated transcription. Ubiquitinated. Polyubiquitination however does not lead to its degradation.

Similarity:

Belongs to the histone deacetylase family. HD type 2 subfamily.

SWISS:

Q9UQL6

Gene ID:

10014

Database links:

[Entrez Gene: 10014](#) Human

[Entrez Gene: 15184](#) Mouse

[Entrez Gene: 84580](#) Rat

[Omim: 605315](#) Human


[SwissProt: Q9UQL6](#) Human

[SwissProt: Q9Z2V6](#) Mouse

[Unigene: 438782](#) Human

[Unigene: 22665](#) Mouse

**Product
Picture**



Sample:

HeLa(Human) Cell Lysate at 30 ug


293T(Human) Cell Lysate at 30 ug

Primary: Anti- Phospho-HDAC5 (Ser259) (SL10326R) at 1/1000 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 119 kD


Observed band size: 119 kD



Paraformaldehyde-fixed, paraffin embedded (mouse colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-HDAC5 (Ser259)) Polyclonal Antibody, Unconjugated (SL10326R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Phospho-HDAC5 (Ser259)) Polyclonal Antibody, Unconjugated (SL10326R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (1M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-Phospho-HDAC5 (Ser259) Polyclonal Antibody,
Unconjugated(SL10326R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining