

Rabbit Anti-Tenascin C antibody

SL1039R

Product Name Tenascin C

Chinese Name 细胞粘合素(固生蛋白)抗体

Alias Tn-C; Cytotactin; Glioma associated extracellular matrix antigen; GMEM; GP 150 225; Hexabrachion; HXB ; JI; Myotendinous antigen; Neuronectin; Tenascin C; TenascinC; TN; TN C; TNC; TENA_HUMAN.

Research Area Tumour immunology

Immunogen Species Rabbit

Clonality Polyclonal

React Species Human, Mouse, Rat, (predicted: Pig,)

Applications IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 (Paraffin sections need antigen repair)
not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.

Theoretical molecular weight 239kDa

Cellular localization Extracellular matrix Secretory protein

Form Liquid

Concentration 1mg/ml

immunogen KLH conjugated synthetic peptide derived from human Tn-C: 1001-1100/2199

Lsotype IgG

Purification affinity purified by Protein A

Buffer Solution 1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.

Storage Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

Attention This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

PubMed [PubMed](#)

Tenascin, also known as hexabrachion and cytotactin, is an extracellular matrix protein with a spatially and temporally restricted tissue distribution. It is a hexameric, multidomain protein with disulfide linked subunits of 190 to 240 kD, originally characterized as 'myotendinous antigen.' In the embryo it is present in dense mesenchyme surrounding developing epithelia and in developing cartilage and bone. In the adult, tenascin remains present in tendons and myotendinous junctions in the perichondrium and periosteum, as well as in smooth muscle.

Function:

Extracellular matrix protein implicated in guidance of migrating neurons as well as axons during development, synaptic plasticity as well as neuronal regeneration. Promotes neurite outgrowth from cortical neurons grown on a monolayer of astrocytes. Ligand for integrins alpha-8/beta-1, alpha-9/beta-1, alpha-V/beta-3 and alpha-V/beta-6.

Subunit:

Homohexamer; disulfide-linked. A homotrimer may be formed in the triple coiled-coil region and may be stabilized by disulfide rings at both ends. Two of such half-hexabrachions may be disulfide linked within the central globule. Interacts with CSPG4.

**Product
Detail**

Subcellular Location:

Secreted, extracellular space, extracellular matrix.

Similarity:

Belongs to the tenascin family.
Contains 15 EGF-like domains.
Contains 1 fibrinogen C-terminal domain.
Contains 15 fibronectin type-III domains.

SWISS:

P24821

Gene ID:

3371

Database links:

[Entrez Gene: 3371](#) Human

[Entrez Gene: 21923](#) Mouse

[Entrez Gene: 116640](#) Rat

[Omim: 187380](#) Human

[SwissProt: P24821](#) Human

[SwissProt: Q80YX1](#) Mouse

[SwissProt: B2LYI9](#) Rat

[Unigene: 143250](#) Human


[Unigene: 454219](#) Mouse

[Unigene: 980](#) Mouse

[Unigene: 12723](#) Rat

Tenascin-C(TN-C)是重要的 Extracellular matrix(ECM)成分之一,在组织器官的发生发育中发挥着极其重要的作用;其表达具有复杂的时空分布,并受多种因素(cell factor、激素等)调节.目前主要用于消化系统 Tumour 方面的研究。

**Product
Picture**




Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (1M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Tenascin C/Tn-C Polyclonal Antibody,

Unconjugated(SL1039R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: mouse stomach tissue; 4% Paraformaldehyde-fixed and

paraffin-embedded;

Antigen retrieval: citrate buffer (1M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Tenascin C/Tn-C Polyclonal Antibody,

Unconjugated(SL1039R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining