SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Marker Primary Antibody]

Rabbit Anti-BACE2/Biotin Conjugated antibody

Catalog Number: SL10494R-Bio

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	Anti-BACE2/Biotin
Immunogen:	KLH conjugated synthetic peptide derived from human BACE2
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Applications:	IHC-P=1:100-500 IHC-F=1:100-500 ICC/IF=1:50-200 IF=1:100-500 ELISA=1:500-5000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	(predicted:Human,Mouse,Rat,Dog,Pig,Cow,Sheep)
Background:	background: Cerebral deposition of amyloid beta peptide is an early and critical feature of Alzheimer's disease and a frequent complication of Down syndrome. Amyloid beta peptide is generated by proteolytic cleavage of amyloid precursor protein by 2 proteases, one of which is the protein encoded by this gene. This gene localizes to the 'Down critical region' of chromosome 21. The encoded protein, a member of the peptidase A1 protein family, is a type I integral membrane glycoprotein and aspartic protease. Three transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]. **Function:** Responsible for the proteolytic processing of the amyloid precursor protein (APP). Cleaves APP, between residues 690 and 691, leading to the

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

	generation and extracellular release of beta-cleaved soluble APP, and a corresponding cell-associated C-terminal fragment which is later released by gamma-secretase. It has also been shown that it can cleave APP between residues 671 and 672. **Subunit:** Monomer. Interacts ith RTN3 and RTN4. **Subcellular Location:** Membrane. Golgi apparatus. Endoplasmic reticulum. Endosome. Cell surface. **Tissue Specificity:** Brain. Present in neurons within the hippocampus, frontal cortex and temporal cortex (at protein level). Expressed at low levels in most peripheral tissues and at higher levels in colon, kidney, pancreas, placenta, prostate, stomach and trachea. Expressed at low levels in the brain. Found in spinal cord, medulla oblongata, substantia nigra and locus coruleus. Expressed in the ductal epithelium of both normal and malignant prostate. **Post-translational modifications:** Undergoes autoproteolytic cleavage. Glycosylated. **Similarity:** Belongs to the peptidase A1 family. **Database links:** **Entrez Gene:** 25825 Human **Entrez Gene:** 56175 Mouse **Entrez Gene:** 288227 Rat Omim: 605668 Human **SwissProt:** Q9Y5Z0 Human **SwissProt:** Q9JL18 Mouse **SwissProt:** Q6IE75 Rat **Unigene:** 529408 Human **Unigene:** 97885 Mouse **Unigene:** 38581 Rat Important **Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.