

## Rabbit Anti-SLITRK5 antibody

SL11959R

<b>Product Name</b>	SLITRK5
<b>Chinese Name</b>	神经突触相关蛋白 SLITRK5 抗体
<b>Alias</b>	bA364G4.2; KIAA0918; Leucine rich repeat containing 11; Leucine rich repeat containing protein 11; LRRC 11; LRRC11; SLIT and NTRK like family member 5; SLIT and NTRK like protein 5; Slit and trk like gene 5; SLITRK 5; SLIK5_HUMAN.
<b>Research Area</b>	Cell biology Neurobiology The cell membrane 蛋白
<b>Immunogen Species</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>React Species</b>	Human(predicted:Mouse,Rat,Dog,Pig,Cow,Horse,Sheep) WB=1:500-2000 (Paraffin sections need antigen repair)
<b>Applications</b>	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Theoretical molecular weight</b>	103kDa
<b>Cellular localization</b>	The cell membrane
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>immunogen</b>	KLH conjugated synthetic peptide derived from human SLITRK5: 301-400/958 <Extracellular>
<b>Lsotype</b>	IgG
<b>Purification</b>	affinity purified by Protein A
<b>Buffer Solution</b>	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
<b>Storage</b>	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
<b>Attention</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>PubMed</b>	<a href="#">PubMed</a>
<b>Product Detail</b>	The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a

hydrophobic  $\alpha$ / $\beta$  horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. SLITRK5 (SLIT and NTRK-like family, member 5), also known as LRRC11 (leucine-rich repeat-containing protein 11), is a 958 amino acid single-pass type I membrane protein that contains 16 LRR repeats and belongs to the SLITRK family. Expressed at high levels in the cerebral cortex, but also present in areas of the spinal cord and medulla, SLITRK5 functions to suppress neurite outgrowth, thereby playing a regulatory role in neuronal function. The gene encoding SLITRK5 maps to human chromosome 13, which houses over 400 genes, such as BRCA2 and RB1, and comprises nearly 4% of the human genome. Trisomy 13, also known as Patau syndrome, is deadly and the few who survive past one year suffer from permanent neurologic defects, difficulty eating and vulnerability to serious respiratory infections.

**Function:**

SLIT and NTRK-like family 5 (SLITRK5) is a member a protein family consisting of six homologous transmembrane proteins (SLITRK1-6) that share two conserved leucine-rich repeat domains in the extracellular domain and have significant homology to Slit, a secreted axonal growth-controlling protein. These proteins are also homologous to trk neurotrophin receptors in their intracellular domains. Expression of SLITRK proteins is highly restricted to neural and brain tumor tissues, but varies within the protein family. Like every other SLITRK protein except SLITRK1, overexpression of SLITRK5 inhibited neurite outgrowth in cultured neurons, suggesting that these proteins are involved in the control of neurite outgrowth.

**Subcellular Location:**

Membrane; Single-pass type I membrane protein.

**Tissue Specificity:**

Expressed predominantly in the cerebral cortex of the brain but also at low levels in the spinal cord and medulla.

**Similarity:**

Belongs to the SLITRK family.  
Contains 12 LRR (leucine-rich) repeats.  
Contains 2 LRRCT domains.  
Contains 1 LRRNT domain.

**SWISS:**

O94991

**Gene ID:**  
26050

**Database links:**

[Entrez Gene: 26050](#) Human

[Entrez Gene: 75409](#) Mouse

[Entrez Gene: 306152](#) Rat


[Omim: 609680](#) Human

[SwissProt: O94991](#) Human

[SwissProt: Q810B7](#) Mouse

[Unigene: 591208](#) Human

**Product Picture**





A549(Human) Cell Lysate at 30 ug

Primary: Anti- SLITRK5 (SL11959R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 103 kD

Observed band size: 100 kD