SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Marker Primary Antibody]

Rabbit Anti-Leupaxin/APC Conjugated antibody

Catalog Number: SL13653R-APC

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	Anti-Leupaxin/APC
Immunogen:	KLH conjugated synthetic peptide derived from human Leupaxin
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Applications:	IF=1:50-200 ICC/IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	(predicted:Human,Mouse,Rat,Dog,Pig,Horse)
Background:	background: Leupaxin is a 386 amino acid cytoplasmic protein and member of the paxillin family. Leupaxin is highly expressed in lymphoid tissues such as spleen, lymph node, thymus and appendix, with low expression in bone marrow and fetal liver. Consisting of four leucine-rich LD-motifs at the N-terminus and four LIM domains at the C-terminus, leupaxin associates with a member of the focal adhesion kinase family, PYK2, in lymphoid cells. The leupaxin and PYK2 complex is involved in cell type-specific signaling in which it regulates signaling at sites of adhesion. Leupaxin is a substrate for tyrosine kinase in lymphoid cells and is suggested to participate in and be regulated by tyrosine kinase activity. Leupaxin may be a potential progression marker for a subset of prostate cancer and may act as a novel coactivator of the androgen receptor. **Function:** Leupaxin is preferentially expressed in hematopoietic cells and is most homologous to the focal adhesion protein, paxillin. It may function in cell

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

	type specific signaling by associating with PYK2, a member of focal adhesion kinase family. As a substrate for a tyrosine kinase in lymphoid cells, this protein may also function in, and be regulated by tyrosine kinase activity. **Subcellular Location:** Cytoplasm **Database links:** **Entrez Gene:** 9404 Human Omim: 605390 Human **SwissProt:** O60711 Human **Unigene:** 125474 Human Important **Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.