SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Marker Primary Antibody]

Rabbit Anti-ELK1/AF647 Conjugated antibody

Catalog Number: SL1398R-AF647

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	Anti-ELK1/AF647
Immunogen:	KLH conjugated synthetic peptide derived from human ELK1
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Applications:	Flow-Cyt=1ug/Test IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Human,Mouse,Rat(predicted:Cow,Horse)
Background:	background: This gene is a member of the Ets family of transcription factors and of the ternary complex factor (TCF) subfamily. Proteins of the TCF subfamily form a ternary complex by binding to the the serum response factor and the serum response element in the promoter of the c-fos proto-oncogene. The protein encoded by this gene is a nuclear target for the ras-raf-MAPK signaling cascade. This gene produces multiple isoforms by using alternative translational start codons and by alternative splicing. Related pseudogenes have been identified on chromosomes 7 and 14. [provided by RefSeq, Mar 2012]. **Function:** Stimulates transcription. Binds to purine-rich DNA sequences. Can form a ternary complex with the serum response factor and the ETS and SRF motifs of the fos serum response element. **Subunit:** Interacts in its sumoylated form with PIAS2/PIASX which enhances its transcriptional activator activity. Interacts with MAD2L2; the interaction is

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

	direct and promotes phosphorylation by the kinases MAPK8 and/or MAPK9. **Subcellular Location:** Nucleus. **Tissue Specificity:** Lung and testis. **Post-translational modifications:** Sumoylation represses transcriptional activator activity as it results in recruitment of HDAC2 to target gene promoters which leads to decreased histone acetylation and reduced transactivator activity. It also regulates nuclear retention. On mitogenic stimulation, phosphorylated on C-terminal serine and threonine residues by MAPK1. Ser-383 and Ser-389 are the preferred sites for MAPK1. In vitro, phosphorylation by MAPK1 potentiates ternary complex formation with the serum responses factors, SRE and SRF. Also phosphorylated on Ser-383 by MAPK8 and/or MAKP9. Phosphorylation leads to loss of sumoylation and restores transcriptional activator activity. Phosphorylated and activated by CAMK4, MAPK11, MAPK12 and MAPK14. **Similarity:** Belongs to the ETS family. Contains 1 ETS DNA-binding domain. **Database links:** **Entrez Gene:** 2002 Human **Entrez Gene:** 13712 Mouse **Entrez Gene:** 314436 Rat Omim: 311040 Human **SwissProt:** P19419 Human **SwissProt:** P41969 Mouse **Unigene:** 181128 Human **Unigene:** 405823 Mouse **Unigene:** 30029 Rat Important **Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. transcriptional regulatory factor（Transcriptin Regulators） The cellular transcription factor ELK1 is a member of the ETS oncogene family.