SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Rabbit Anti-RBBP6 antibody

Catalog Number: SL1483R

FOR RESEARCH USE ONLY

[DATASHEET]


Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	RBBP6
Immunogen:	KLH conjugated synthetic peptide derived from human RBBP6: 2-150/1792
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	IHC-P=1:100-500 IHC-F=1:100-500 IF=1:100-500 （Paraffin sections need antigen repair） not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Mouse(predicted:Human,Rat)
Predicted MW.:	201kDa
Subcellular Location:	The nucleus cytoplasmic
Background:	he retinoblastoma tumor suppressor (pRB) protein binds with many other proteins. In various human cancers, pRB suppresses cellular proliferation and is inactivated. Cell cycle-dependent phosphorylation regulates the activity of pRB. This gene encodes a protein which binds to underphosphorylated but not phosphorylated pRB. Multiple alternatively spliced transcript variants that encode different isoforms have been found for this gene. [provided by RefSeq, Jul 2008] **Function:** E3 ubiquitin-protein ligase which promotes ubiquitination of YBX1, leading to its degradation by the proteasome. May play a role as a scaffold protein to promote the assembly of the p53/TP53-MDM2 complex, resulting in increase of MDM2-mediated ubiquitination and degradation of p53/TP53; may function as negative regulator of p53/TP53, leading to both apoptosis and cell growth.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

	Subunit: Interacts with p53/TP53 and RB1. Interacts also with MDM2 and YBX1. Interacts with NEK6. **Subcellular Location:** Nucleus, nucleolus. Chromosome. Cytoplasm, cytoskeleton, centrosome. Note=Colocalizes with mitotic chromosomes. Co-localizes with NEK6 in the centrosome. **Tissue Specificity:** Highly expressed in the placenta and testis. Expressed at lower levels in the brain, heart, kidney, liver and lung. Overexpressed in esophageal cancer. **Post-translational modifications:** Phosphorylated by NEK6. **Similarity:** Contains 1 CCHC-type zinc finger. Contains 1 DWNN domain. Contains 1 RING-type zinc finger. **Database links:** **Entrez Gene:** 5930 Human **Entrez Gene:** 19647 Mouse **Entrez Gene:** 308968 Rat Omim: 600938 Human **SwissProt:** Q7Z6E9 Human **SwissProt:** P97868 Mouse **Unigene:** 188553 Human **Unigene:** 4480 Mouse Signaling Intermediates P2P-P, also known as PACT, i.e., P53 associated cellular protein-testes derived. P2P-P protein is a negative regulator of p53 and plays an important role in cell growth and embryonic development, cell differentiation, pre-mRNA splicing, cell cycle arrest, and apoptosis.

[VALIDATION IMAGES]



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RBBP6) Polyclonal Antibody, Unconjugated (SL1483R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.