SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Rabbit Anti-C1orf33 antibody

Catalog Number: SL15064R

FOR RESEARCH USE ONLY

[DATASHEET]


Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	C1orf33
Immunogen:	KLH conjugated synthetic peptide derived from human C1orf33: 1-80/239
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	IHC-P=1:100-500 IHC-F=1:100-500 IF=1:100-500 （Paraffin sections need antigen repair） not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Human(predicted:Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep)
Predicted MW.:	28kDa
Subcellular Location:	The nucleus cytoplasmic
Background:	This gene encodes a protein sharing a low level of sequence similarity with ribosomal protein P0. While the precise function of the encoded protein is currently unknown, it appears to be involved in mRNA turnover and ribosome assembly. [provided by RefSeq, Jul 2008] **Function:** Involved in mRNA turnover and ribosome assembly (By similarity). **Subcellular Location:** Nucleus, nucleolus. **Similarity:** Belongs to the ribosomal protein L10P family. **Database links:** **Entrez Gene:** 509932 Cow **Entrez Gene:** 51154 Human **Entrez Gene:** 69902 Mouse GenBank: NP_057267 Human

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

	SwissProt: A4FV84 Cow **SwissProt:** Q9UKD2 Human **SwissProt:** Q9D0I8 Mouse **Unigene:** 463797 Human **Unigene:** 27871 Mouse **Unigene:** 296860 Mouse **Unigene:** 441201 Mouse

[VALIDATION IMAGES]



Paraformaldehyde-fixed, paraffin embedded (human colon cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (C1orf33) Polyclonal Antibody, Unconjugated (SL15064R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.