SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Rabbit Anti-GALNT4 antibody

Catalog Number: SL16220R

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	GALNT4
Immunogen:	KLH conjugated synthetic peptide derived from human GALNT4: 51-150/578
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	IHC-P=1:100-500 IHC-F=1:100-500 ICC/IF=1:100-500 IF=1:100-500 ELISA=1:5000-10000 （Paraffin sections need antigen repair） not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	(predicted: Human, Mouse, Rat, Cow, Horse, Rabbit, Sheep)
Predicted MW.:	67kDa
Subcellular Location:	cytoplasmic
Background:	This gene encodes a member of the UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes. GalNAc-Ts initiate mucin-type O-linked glycosylation in the Golgi apparatus by catalyzing the transfer of GalNAc to serine and threonine residues on target proteins. They are characterized by an N-terminal transmembrane domain, a stem region, a lumenal catalytic domain containing a GT1 motif and Gal/GalNAc transferase motif, and a C-terminal ricin/lectin-like domain. GalNAc-Ts have different, but overlapping, substrate specificities and patterns of expression. In vitro, the encoded protein can complement other GalNAc-Ts in the complete O-glycosylation of the mucin-1 tandem repeat and can O-

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

	glycosylate the P-selectin glycoprotein ligand-1 molecule. The coding region of this gene is contained within a single exon. Fusion transcripts, which combine part of this gene with the 5' exons of the neighboring POC1B (POC1 centriolar protein homolog B) gene, also exist. [provided by RefSeq, Dec 2010] **Function:** Catalyzes the initial reaction in O-linked oligosaccharide biosynthesis, the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. Has a highest activity toward Muc7, EA2 and Muc2, with a lowest activity than GALNT2. Glycosylates 'Thr-57' of SELPLG. **Subcellular Location:** Golgi apparatus membrane. **Tissue Specificity:** Ubiquitous. Highly expressed in mucous cells. **Similarity:** Belongs to the glycosyltransferase 2 family. GalNAc-T subfamily. Contains 1 ricin B-type lectin domain. **Database links:** **Entrez Gene:** 8693 Human **Entrez Gene:** 14426 Mouse **Entrez Gene:** 500826 Rat Omim: 603565 Human **SwissProt:** Q8N4A0 Human **SwissProt:** O08832 Mouse **Unigene:** 25130 Human **Unigene:** 314 Mouse

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.