

## Rabbit Anti-CCR10/AF350 Conjugated antibody

SL1710R-AF350

<b>Product Name</b>	Anti-CCR10/AF350
<b>Chinese Name</b>	AF350 标记的细胞表面 Chemokine 受体 10 抗体
<b>Alias</b>	C-C chemokine receptor 10; C C chemokine receptor type 10; C C CKR 10; CC chemokine receptor 10; CC CKR 10; CCR 10; CCR-10; CCR10; Chemokine (C C motif) receptor 10; G protein coupled receptor 2; G protein-coupled receptor 2; GPR 2; GPR2.
<b>Research Area</b>	The cell membrane 受体
<b>Immunogen Species</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>React Species</b>	Human,Mouse(predicted:Rat,Dog,Pig,Cow,Rabbit)
<b>Applications</b>	Flow-Cyt=5µg/Test,IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight</b>	40kDa
<b>Form</b>	Lyophilized or Liquid
<b>Concentration</b>	1mg/ml
<b>immunogen</b>	KLH conjugated synthetic peptide derived from human CCR-10 (13-45aa)
<b>Lsotype</b>	IgG
<b>Purification</b>	affinity purified by Protein A
<b>Storage Buffer</b>	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol
<b>Storage</b>	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>Product Detail</b>	<b>background:</b> Chemokines are a group of small (approximately 8 to 14 kD), mostly basic, structurally related molecules that regulate cell trafficking of various types of leukocytes through interactions with a subset of 7-transmembrane, G

protein-coupled receptors. Chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis. Chemokines are divided into 2 major subfamilies, CXC and CC, based on the arrangement of the first 2 of the 4 conserved cysteine residues; the 2 cysteines are separated by a single amino acid in CXC chemokines and are adjacent in CC chemokines. CCR10 is the receptor for CCL27 (SCYA27; MIM 604833); CCR10-CCL27 interactions are involved in T cell-mediated skin inflammation (Homey et al., 2002 [PubMed 11821900]).[supplied by OMIM, Mar 2008]

**Function:**

Receptor for chemokines SCYA27 and SCYA28. Subsequently transduces a signal by increasing the intracellular calcium ions level and stimulates chemotaxis in a pre-B cell line.

**Subcellular Location:**

Cell membrane; Multi-pass membrane protein.

**Tissue Specificity:**

Expressed at high levels in adult testis, small intestine, fetal lung, fetal kidney. Weaker expression was observed in many other adult tissues including spleen, thymus, lymph node, Peyer patches, colon, heart, ovary, peripheral blood lymphocytes, thyroid and spinal cord. Also expressed by melanocytes, dermal fibroblasts, dermal microvascular endothelial cells. Also detected in T-cells and in skin-derived Langerhans cells.

**Similarity:**

Belongs to the G-protein coupled receptor 1 family.

**Database links:**

[Entrez Gene: 2826](#) Human

[Entrez Gene: 12777](#) Mouse

[Omim: 600240](#) Human

[SwissProt: P46092](#) Human

[SwissProt: Q9JL21](#) Mouse

[Unigene: 278446](#) Human

[Unigene: 103813](#) Mouse

[Unigene: 8021](#) Mouse



SunLong Biotech Co.,LTD  
Tel: 0086-571-56623320 Fax:0086-571-56623318  
E-mail:sales@sunlongbiotech.com  
www.sunlongbiotech.com

---

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.