SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Marker Primary Antibody]

Rabbit Anti-phospho-MCAK (Ser95)/FITC Conjugated antibody

Catalog Number: SL18714R-FITC

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	Anti-phospho-MCAK (Ser95)/FITC
Immunogen:	KLH conjugated synthetic peptide derived from human phospho-MCAK (Ser95)
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Applications:	IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Human(predicted:Horse,Rabbit)
Background:	background: This gene encodes a kinesin-like protein that functions as a microtubule-dependent molecular motor. The encoded protein can depolymerize microtubules at the plus end, thereby promoting mitotic chromosome segregation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2014] **Function:** Promotes ATP-dependent removal of tubulin dimers from microtubules. Regulates the turnover of microtubules at the kinetochore and functions in chromosome segregation during mitosis. **Subcellular Location:** Cytoplasm > cytoskeleton. Nucleus. Chromosome > centromere. Chromosome > centromere > kinetochore. Associates with the microtubule network at the growing distal tip (the plus-end) of microtubules, probably through interaction with MTUS2/TIP150 and MAPRE1 (By similarity). Centromeric localization requires the presence of BUB1 and SGOL2.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

	Tissue Specificity: Expressed at high levels in thymus and testis, at low levels in small intestine, the mucosal lining of colon, and placenta, and at very low levels in spleen and ovary; expression is not detected in prostate, peripheral blood Leukocytes, heart, brain, lung, liver, skeletal muscle, kidney or pancreas. Isoform 2 is testis-specific. **Post-translational modifications:** Phosphorylated upon DNA damage, probably by ATM or ATR. Phosphorylation by STK12, regulates association with centromeres and kinetochores and the microtubule depolymerization activity. **Similarity:** Belongs to the kinesin-like protein family. MCAK/KIF2 subfamily. Contains 1 kinesin-motor domain. **Database links:** **Entrez Gene:** 11004 Human **Entrez Gene:** 73804 Mouse Omim: 604538 Human **SwissProt:** Q99661 Human **SwissProt:** Q922S8 Mouse **Unigene:** 720061 Human **Unigene:** 247651 Mouse Important **Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.