SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Rabbit Anti-VANGL2 antibody

Catalog Number: SL23066R

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	VANGL2
Immunogen:	KLH conjugated synthetic peptide derived from human VANGL2: 151-250/521 <Extracellular>
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB=1:500-2000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Human,Mouse(predicted:Rat,Pig,Cow,Horse,Sheep)
Predicted MW.:	60kDa
Subcellular Location:	The cell membrane
Background:	The Vang family of proteins are integral membrane proteins that are homologs of the Drosophila tissue polarity gene strabismus. The gene encoding for Van Gogh-like protein 1 (Vangl1), also designated Strabismus 2 (STB2), localizes to chromosome 1p11-p13.1. Van Gogh-like protein 2 (Vangl2), also designated Strabismus1 (STB1), localizes on chromosome 1q22-q23. Vangl1 is expressed in testis and ovary, but also in gastric and pancreatic cancer. Vangl proteins play a key developmental role in establishing planar cell polarity (PCP) and in regulating convergent extension (CE) movements during embryogenesis. Vangl1 and Vangl2 are both down-regulated in several cancer cell lines and primary tumors. Involved in the control of early morphogenesis and patterning of both axial midline structures and the development of neural plate. Plays a role in the regulation of planar cell polarity, particularly in the orientation of stereociliary bundles in the cochlea. Required for polarization and movement of

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

	myocardializing cells in the outflow tract and seems to act via RHOA signaling to regulate this process. **Function:** Involved in the control of early morphogenesis and patterning of both axial midline structures and the development of neural plate. Plays a role in the regulation of planar cell polarity, particularly in the orientation of stereociliary bundles in the cochlea. Required for polarization and movement of myocardializing cells in the outflow tract and seems to act via RHOA signaling to regulate this process (By similarity). **Subunit:** Interacts through its C-terminal region with the N-terminal half of DVL1, DVL2 and DVL3. The PDZ domain of DVL1, DVL2 and DVL3 is required for the interaction. Also interacts with the PDZ domains of MAGI3, SCRIB/SCRB1 and FZD3 (By similarity). **Subcellular Location:** Membrane; Multi-pass membrane protein (Potential). DISEASE: Defects in VANGL2 are a cause of neural tube defects (NTD) [MIM:182940]. NTD are congenital malformations of the central nervous system and adjacent structures related to defective neural tube closure during the first trimester of pregnancy. Failure of neural tube closure can occur at any level of the embryonic axis. Common NTD forms include anencephaly, myelomeningocele and spina bifida, which result from the failure of fusion in the cranial and spinal region of the neural tube. NTDs have a multifactorial etiology encompassing both genetic and environmental components. **Similarity:** Belongs to the Vang family. **Database links:** **Entrez Gene:** 57216 Human **Entrez Gene:** 93840 Mouse **Entrez Gene:** 289229 Rat Omim: 600533 Human **SwissProt:** Q9ULK5 Human **SwissProt:** Q91ZD4 Mouse **SwissProt:** P84889 Rat **Unigene:** 99477 Human **Unigene:** 36148 Mouse **Unigene:** 392110 Mouse **Unigene:** 198958 Rat

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

[VALIDATION IMAGES]



Sample:
NIH/3T3(Mouse) Cell Lysate at 30 ug
Primary: Anti-VANGL2 (SL23066R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 60 kD
Observed band size: 60 kD

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.