

Rabbit Anti-Caspase-2 p13/Cy5.5 Conjugated antibody

SL23456R-Cy5. 5

Product Name	Anti-Caspase-2 p13/Cy5.5
Chinese Name	Cy5.5 标记的活化半胱氨酸蛋白酶蛋白-2 抗体 Caspase-2 subunit p13; CASP 2; CASP-2; Casp2; CASP2_HUMAN; Caspase 2; Caspase 2 apoptosis related cysteine peptidase; Caspase2; ICH 1; ICH 1 protease; ICH 1L; ICH1; ICH1 protease; ICH1L; NEDD-2; NEDD2; Neural precursor cell expressed developmentally down-regulated protein 2; PPP1R57; Protease ICH-1; Protein phosphatase 1 regulatory subunit 57.
Alias	
Research Area	Tumour Cell biology Signal transduction Apoptosis The new supersedes the old
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Human,Rat(predicted:Mouse,Pig,Horse,Rabbit,Sheep) IF=1:100-500
Applications	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight	13/51kDa
Form	Lyophilized or Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human Caspase-2 subunit p13
Lsotype	IgG
Purification	affinity purified by Protein A
Storage Buffer	Preservative: 15mM Sodium Azide, Constituents: 1% BSA, 1M PBS, pH 7.4. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Storage	
Product Detail	background: This gene encodes a member of the cysteine-aspartic acid protease (caspase)

family. Caspases mediate cellular apoptosis through the proteolytic cleavage of specific protein substrates. The encoded protein may function in stress-induced cell death pathways, cell cycle maintenance, and the suppression of tumorigenesis. Increased expression of this gene may play a role in neurodegenerative disorders including Alzheimer's disease, Huntington's disease and temporal lobe epilepsy. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2011]

Function:

Involved in the activation cascade of caspases responsible for apoptosis execution. Might function by either activating some proteins required for cell death or inactivating proteins necessary for cell survival.

Subunit:

Heterotetramer that consists of two anti-parallel arranged heterodimers, each one formed by a p18 subunit and a p12 subunit. Interacts with LRDD. Interacts with NOL3 (via CARD domain); inhibits CASP2 activity in a phosphorylation-dependent manner.

Tissue Specificity:

Expressed at higher levels in the embryonic lung, liver and kidney than in the heart and brain. In adults, higher level expression is seen in the placenta, lung, kidney, and pancreas than in the heart, brain, liver and skeletal muscle.

Post-translational modifications:

The mature protease can process its own propeptide, but not that of other caspases.

Similarity:

Belongs to the peptidase C14A family.
Contains 1 CARD domain.

Database links:

[Entrez Gene: 835](#) Human

[Entrez Gene: 12366](#) Mouse

[Entrez Gene: 64314](#) Rat

[Omim: 600639](#) Human

[SwissProt: P42575](#) Human



[SwissProt: P29594](#) Mouse

[SwissProt: P55215](#) Rat

[Unigene: 368982](#) Human

[Unigene: 3921](#) Mouse

[Unigene: 1438](#) Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.