SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

**[ Primary Antibody ]**

# Rabbit Anti-RGC32 antibody

**Catalog Number: SL23862R**

**FOR RESEARCH USE ONLY**

**[ DATASHEET ]**


| | |
|---|---|
| **Host:** | Rabbit |
| **Isotype:** | IgG |
| **Clonality:** | Polyclonal |
| **Target:** | RGC32 |
| **Immunogen:** | KLH conjugated synthetic peptide derived from human RGC32: 31-100/137 |
| **Purification:** | affinity purified by Protein A |
| **Concentration:** | 1mg/1ml |
| **Storage:** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. |
| **Applications:** | IHC-P=1:100-500<br>IHC-F=1:100-500<br>IF=1:100-500 （Paraffin sections need antigen repair）<br>not yet tested in other applications.<br>optimal dilutions/concentrations should be determined by the end user. |
| **Reactivity:** | Rat(predicted:Human,Cow,Horse) |
| **Predicted MW.:** | 15kDa |
| **Subcellular Location:** | The nucleus cytoplasmic |
| **Background:** | RGC32, also known as C13orf15, is a 137 amino acid protein that localizes to the cytoplasm, as well as to the nucleus and the centrosome. Expressed at high levels in kidney, pancreas and skeletal muscle and at lower levels in brain, heart and placenta, RGC32 functions to modulate the activity of cell cycle-specific kinases, thereby regulating cell cycle progression. Additionally, RGC32 may promote cell cycle arrest at the G2/M phase transition and is thought to inhibit the growth of glioma cells, possibly functioning as a tumor suppressor.<br>Conversely, overexpression of RGC32 may promote cell replication and assist in the pathogenesis of malignancies, suggesting that RGC32 also participates in tumor transformation and progression. RGC32 activity is induced by complement activation and by p53 in response to DNA damage. Multiple isoforms of RGC32 exist as a result of alternative splicing events. |

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

| | |
|---|---|
| | **Function:** Modulates the activity of cell cycle-specific kinases. Enhances CDK1 activity. May contribute to the regulation of the cell cycle. May inhibit growth of glioma cells by promoting arrest of mitotic progression at the G2/M transition. Fibrogenic factor contributing to the pathogenesis of renal fibrosis through fibroblast activation.<br>**Subunit:** Interacts with SMAD3. Interacts with CDK1 and PLK1.<br>**Subcellular Location:** Cytoplasm. Nucleus.<br>Cytoplasm ?cytoskeleton ?centrosome.<br>**Note:** Cytoplasmic in unstimulated cells. Nuclear after activation by complement. Associated with the centrosome during prometaphase and metaphase.<br>**Tissue Specificity:** Detected in brain, heart and liver (at protein level). Highly expressed in liver, skeletal muscle, kidney and pancreas. Detected at lower levels in heart, brain and placenta. Detected in aorta endothelial cells. Overexpressed in colon, breast, prostate, bladder, lung, and ovarian cancer tissues.<br>**Database links:**<br>**Entrez Gene:** 28984 Human<br>Omim: 610077 Human<br>**SwissProt:** Q9H4X1 Human |

## [ VALIDATION IMAGES ]



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RGC32) Polyclonal Antibody, Unconjugated (SL23862R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.