

Rabbit Anti-phospho-FSCN1 (Ser39) antibody

SL2833R

Product Name phospho-FSCN1 (Ser39)

Chinese Name 磷酸化纤维束蛋白同源物 1 抗体

Alias

Fascin (phospho Ser39); Fascin (phospho S39); 55 kDa actin bundling protein; Actin bundling protein; FAN 1; FAN1; Fascin 1; Fascin homolog 1 actin bundling protein (Strongylocentrotus purpuratus); Fascin homolog 1; Fascin1; FLJ38511; FSCN 1; FSCN1; HSN; p55; Singed (Drosophila) like (sea urchin fascin homolog like); Singed drosophila homolog like; Singed like (fascin homolog sea urchin) (Drosophila); Singed like protein; SNL; Strongylocentrotus purpuratus.

Product Type Phosphorylated anti

Research Area Cell biology Signal transduction Binding protein

Immunogen Species Rabbit

Clonality Polyclonal

React Species Human,Mouse,Rat(predicted:Dog,Pig)
WB=1:500-2000

Applications not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.

Theoretical molecular weight 55kDa

Cellular localization cytoplasmic

Form Liquid

Concentration 1mg/ml

immunogen KLH conjugated Synthesised phosphopeptide derived from human FSCN1 around the phosphorylation site of Ser39: AS(p-S)LK

Lsotype IgG

Purification affinity purified by Protein A

Buffer 1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.



Solution

Storage

Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.

Attention

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

PubMed

[PubMed](#)

Human fascin is a highly conserved actin-bundling protein. Fascin, encoded by the human homolog for sn (hsn) gene, has been localized to microspikes and stress fibers of cultured cells where it is thought to be involved in the formation of microfilament bundles. It is expressed predominantly in dendritic cells. Lymphoid cells, myeloid cells and plasma cells are negative. However, Reed Sternberg cells in Hodgkin's lymphoma are positive for fascin staining. Epstein-Barr virus may induce expression of fascin in B cells.

Function:

Organizes filamentous actin into bundles with a minimum of 4.1:1 actin/fascin ratio. Plays a role in the organization of actin filament bundles and the formation of microspikes, membrane ruffles, and stress fibers. Important for the formation of a diverse set of cell protrusions, such as filopodia, and for cell motility and migration.

Subcellular Location:

Cytoplasm > cytoskeleton. Cell projection > filopodium. Cell projection > invadopodium.

**Product
Detail**

Tissue Specificity:

Ubiquitous.

Post-translational modifications:

Phosphorylation on Ser-39 inhibits the actin-binding ability of fascin.

Similarity:

Belongs to the fascin family.

SWISS:

Q16658

Gene ID:

6624

Database links:

[Entrez Gene: 6624](#) Human

[Entrez Gene: 14086](#) Mouse

[Entrez Gene: 683788](#) Rat

[Omim: 602689](#) Human

[SwissProt: Q16658](#) Human

[SwissProt: Q61553](#) Mouse

[SwissProt: P85845](#) Rat


[Unigene: 118400](#) Human

[Unigene: 289707](#) Mouse

[Unigene: 199526](#) Rat

肌动蛋白集束蛋白/圆线虫紫癜抗体。 FSCN1 为 Cytoskeleton 肌动蛋白的一种，该蛋白有蛋白结合,桥连、肌动蛋白丝结合的功能。 主要参与细胞增殖、肌动蛋白 Cytoskeleton 组织和生物发生、肌动蛋白丝束形成。 <

Product
Picture



Sample:

Lane 1: Mouse Cerebrum tissue lysates

Lane 2: Mouse Testis tissue lysates

Lane 3: Mouse Spleen tissue lysates

Lane 4: Rat Cerebrum tissue lysates

Lane 5: Rat Testis tissue lysates

Lane 6: Rat Spleen tissue lysates

Lane 7: Human HepG2 cell lysates

Lane 8: Human K562 cell lysates

Lane 9: Human SH-SY5Y cell lysates

Lane 10: Human U-2os cell lysates

Primary: Anti- phospho-FSCN1 (Ser39) (SL2833R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 55 kDa

Observed band size: 51 kDa