

Rabbit Anti-hnRNP L/Cy5 Conjugated antibody

SL4300R-Cy5

Product Name	Anti-hnRNP L/Cy5
Chinese Name	Cy5 标记的异质核糖核蛋白 L 抗体
Alias	Heterogeneous nuclear ribonucleoprotein L; hnRNP L; hnRNPL; hnRPL; HNRPL_HUMAN; P/OKcl.14; D830027H13Rik; FLJ35509.
Research Area	Cell biology immunology Chromatin and nuclear signals
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Mouse(predicted:Human,Rat,Pig,Cow,Horse,Rabbit,GuineaPig) IF=1:100-500
Applications	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight	65kDa
Form	Lyophilized or Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human hnRNP L
Lsotype	IgG
Purification	affinity purified by Protein A
Storage Buffer	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Product Detail	background: Heterogeneous nuclear RNAs (hnRNAs) which include mRNA precursors and mature mRNAs are associated with specific proteins to form heterogenous ribonucleoprotein (hnRNP) complexes. Heterogeneous nuclear ribonucleoprotein L is among the proteins that are stably associated with hnRNP complexes and along with other hnRNP proteins is likely to play a

major role in the formation, packaging, processing, and function of mRNA. Heterogeneous nuclear ribonucleoprotein L is present in the nucleoplasm as part of the HNRP complex. HNRP proteins have also been identified outside of the nucleoplasm. Exchange of hnRNP for mRNA-binding proteins accompanies transport of mRNA from the nucleus to the cytoplasm. Since HNRP proteins have been shown to shuttle between the nucleus and the cytoplasm, it is possible that they also have cytoplasmic functions. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq].

Subunit:

Identified in a mRNP granule complex, at least composed of ACTB, ACTN4, DHX9, ERG, HNRNPA1, HNRNPA2B1, HNRNPAB, HNRNPD, HNRNPL, HNRNPR, HNRNPU, HSPA1, HSPA8, IGF2BP1, ILF2, ILF3, NCBP1, NCL, PABPC1, PABPC4, PABPN1, RPLP0, RPS3, RPS3A, RPS4X, RPS8, RPS9, SYNCRIP, TROVE2, YBX1 and untranslated mRNAs. Interacts with HNRPLL. Interacts with APEX1; the interaction is DNA-dependent.

Subcellular Location:

Nucleus, nucleoplasm. Cytoplasm.

Similarity:

Contains 3 RRM (RNA recognition motif) domains.

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.