


Rabbit Anti-Mono-Methyl-Histone H3 (Lys23)antibody

SL60104R

Product Name	Mono-Methyl-Histone H3 (Lys23)
Chinese Name	
Product Type	Methylated anti
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Human,Mouse,Rat
Applications	WB=1:500-2000,IP=1:20-100,ChIP=6 ug/5x10 ⁶ cells not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Theoretical molecular weight	17kDa
Cellular localization	The nucleus
Form	Liquid
Concentration	1mg/ml
Lsotype	IgG
Purification	Antigen affinity purification
Buffer Solution	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
PubMed	PubMed
Product Detail	组蛋白翻译后修饰（PTMs）是 Epigenetics 调控染色质结构的关键机制，被称为“组蛋白密码”。组蛋白上的翻译后修饰包括乙酰化，甲基化，磷酸化和近年发现的一些新型酰化修饰。这些组蛋白修饰直接影响染色质和转录因子或其他表观调控子的结合，改变基因组的稳定性和基因转录等。组蛋白甲基化通常发生在核心组蛋白的赖氨酸和精氨酸残基。组蛋

白甲基化既可以促进也可以抑制基因转录，这取决于组蛋白甲基化是发生在赖氨酸还是发生在精氨酸上以及甲基化基团的个数（赖氨酸可以发生一、二、三甲基化，精氨酸可发生单、对称与非对称二甲化）。组蛋白赖氨酸甲基化通常发生在组蛋白 H3 第 4, 9, 27, 36, 79 和组蛋白 H4 的第 20 位赖氨酸上；精氨酸甲基化通常发生在组蛋白 H3 的第 2, 8, 17, 26 和组蛋白 H4 第 3 位精氨酸上。蛋白甲基化酶（HMT）和组蛋白去甲基化酶（HDM）是甲基化修饰主要的调控因子。



Product Picture

Blocking buffer: 5% NFDN/TBST

Primary ab dilution: 1:2000

Primary ab incubation condition: 2 hours at room temperature

Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)

Lysate: HeLa, 3T3, H1299

Protein loading quantity: 20 μ g

Exposure time: 15 s

Predicted MW: 17 kDa



SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

Observed MW: 17 kDa