

Fab'₂ Fragment Goat Anti-Mouse IgG H&L / APC antibody

SL60296G-APC

Product Name	F(ab') ₂ Fragment Goat Anti-Mouse IgG H&L / APC
Chinese Name	APC 标记的(ab) ₂ 片段羊抗小鼠 IgG H&L
Alias	F(ab) ₂ Fragment Goat Anti-Mouse IgG H&L (APC); F(ab) ₂ Fragment Goat Anti-Mouse IgG H&L/APC; F(ab) ₂ Fragment Goat Anti-Mouse IgG(H+L);
Immunogen Species	Goat
Clonality	Polyclonal
React Species	Mouse,
Applications	ICC/IF=1:100-1000,IF=1:100-1000,Flow-Cyt=1:100-1000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Form	Liquid
Concentration	1 mg/ml
immunogen	Native Mouse IgG
Lsotype	IgG
Purification	affinity purified by Protein G
Buffer Solution	10 mM TBS (pH=7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Attention	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. Immunoglobulin G (IgG), is one of the most abundant proteins in serum with normal levels between 8-17 mg/mL in adult blood. IgG is important for our defence against microorganisms and the molecules are produced by B lymphocytes as a part of our adaptive immune response. The IgG molecule has two separate functions; to bind to the pathogen that elicited the response and to recruit other cells and molecules to destroy the antigen. The variability of the IgG pool is generated by somatic recombination and the number of specificities in an individual at a given time point is estimated to be 10 ¹¹
Product Detail	



SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

variants.