

## Rabbit Anti-NRIP/Biotin Conjugated antibody

SL6210R-Bio

<b>Product Name</b>	Anti-NRIP/Biotin
<b>Chinese Name</b>	生物素标记的雄激素受体复合物相关蛋白/核受体相互作用蛋白抗体
<b>Alias</b>	Androgen receptor complex associated protein; ARCAP; IQ motif and WD repeats 1; IQWD1v MSTP055; Nuclear receptor interaction protein; RP4 745I14.1; DCAF6_HUMAN.
<b>Research Area</b>	Tumour Cardiovascular Chromatin and nuclear signals Signal transduction
<b>Immunogen Species</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>React Species</b>	(predicted:Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit)
<b>Applications</b>	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:500-5000 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Molecular weight</b>	96kDa
<b>Form</b>	Lyophilized or Liquid
<b>Concentration</b>	1mg/ml
<b>immunogen</b>	KLH conjugated synthetic peptide derived from human NRIP
<b>Lsotype</b>	IgG
<b>Purification</b>	affinity purified by Protein A
<b>Storage Buffer</b>	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
<b>Storage</b>	
<b>Product Detail</b>	<b>background:</b> WD-repeat proteins are a large family of eukaryotic proteins coordinating multi-protein complex assemblies. Their role has been implicated in multiple cellular processes including signal transduction, transcriptional regulation, cell cycle control and apoptosis. NRIP is a novel 860a.a nuclear protein consisting of seven conserved WD40 domains and one NLS motif. It binds to androgen and glucocorticoid receptors and up-regulates their transcriptional

activity, thereby functioning as a nuclear receptor co-activator. Role of NRIP has been implicated in cell growth and also in cervical and prostate cancer, thus indicating a potential therapeutic intervention. Northern Blot analysis detects a high expression of NRIP in skeletal muscle and testis and low expression in heart, prostate and adrenal gland.

**Function:**

Ligand-dependent coactivator of nuclear receptors. Enhance transcriptional activity of the nuclear receptors NR3C1 and AR. May function as a substrate receptor for CUL4-DDB1 E3 ubiquitin-protein ligase complex.

**Subunit:**

Interacts with the nuclear receptors NR3C1 and AR in the presence of ligand. Interacts with DDB1, CUL4A and CUL4B.

**Subcellular Location:**

Nucleus.

**Tissue Specificity:**

Highly expressed in skeletal muscle and testis. Expressed to a lesser degree in heart, prostate, and adrenal gland.

**Similarity:**

Contains 1 IQ domain.  
Contains 7 WD repeats.

**Database links:**

[Entrez Gene: 55827](#) Human

[Entrez Gene: 74106](#) Mouse

[Entrez Gene: 289181](#) Rat

[Omim: 610494](#) Human

[SwissProt: Q58WW2](#) Human

[SwissProt: Q9DC22](#) Mouse

[Unigene: 435741](#) Human

[Unigene: 227605](#) Mouse

**Important Note:**



SunLong Biotech Co.,LTD  
Tel: 0086-571-56623320 Fax:0086-571-56623318  
E-mail:sales@sunlongbiotech.com  
www.sunlongbiotech.com

---

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.