SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Marker Primary Antibody]

Rabbit Anti-IL-1 Beta , Alexa Fluor® 680 conjugated antibody

Catalog Number: SL6319R-AF680

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	IL-1 Beta, Bodipy Fluor 680 conjugated
Immunogen:	KLH conjugated synthetic peptide derived from human IL-1 Beta: 161-269/269
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20℃ for one year. Avoid repeated freeze/thaw cycles.
Applications:	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Mouse(predicted:Human,Bee)
Predicted MW.:	17/32kDa
Subcellular Location:	Secretory protein
Background:	The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2008]. **Function:** Produced by activated macrophages, IL-1 stimulates thymocyte proliferation by inducing IL-2 release, B-cell maturation and proliferation, and fibroblast growth factor activity. IL-1 proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

	synovial cells. **Subunit:** Monomer. **Subcellular Location:** Secreted. Note=The lack of a specific hydrophobic segment in the precursor sequence suggests that IL-1 is released by damaged cells or is secreted by a mechanism differing from that used for other secretory proteins. **Similarity:** Belongs to the IL-1 family. **Database links:** **Entrez Gene:** 3553 Human **Entrez Gene:** 16176 Mouse **Entrez Gene:** 24494 Rat Omim: 147720 Human **SwissProt:** P01584 Human **SwissProt:** P10749 Mouse **SwissProt:** Q63264 Rat **Unigene:** 126256 Human **Unigene:** 222830 Mouse **Unigene:** 9869 Rat

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.