

Rabbit Anti-GANAB/Biotin Conjugated antibody

SL6720R-Bio

Product Name	Anti-GANAB/Biotin
Chinese Name	生物素标记的 α 葡萄糖苷酶 2 抗体
Alias	Glucosidase II α ; Alpha glucosidase II alpha subunit; Alpha-glucosidase 2; G2AN; GANAB; GANAB_HUMAN; Glu II; Glucosidase alpha neutral AB; Glucosidase II alpha; Glucosidase II subunit alpha; GluII; KIAA0088; Neutral alpha glucosidase AB; Neutral alpha glucosidase AB precursor; Neutral alpha-glucosidase AB.
Research Area	Tumour Cell biology Neurobiology Signal transduction
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Mouse,Rat(predicted:Human,Pig,Cow,Horse,Rabbit,Sheep) WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Applications	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight	104kDa
Form	Lyophilized or Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human GANAB/alpha Glucosidase II
Lsotype	IgG
Purification	affinity purified by Protein A
Storage Buffer	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Storage	
Product Detail	background: Trimming of glucoses from N-linked core glycans on newly synthesized

glycoproteins occurs sequentially through the action of Glucosidases I and II in the endoplasmic reticulum (ER). Glucosidase II is an ER-localized enzyme that contains a and b subunits (Glucosidase IIa and Glucosidase IIb) which form a defined heterodimeric complex. Glucosidase IIa is the catalytic core of the enzyme and can function independently of the b subunit. The sequence of Glucosidase IIb encodes protein rich in glutamic and aspartic acid with a putative ER retention signal (HDEL) at the C-terminus. The phosphorylated form of Glucosidase IIb is localized in the plasma membrane and is highly expressed in FGF-stimulated fibroblasts and epidermal carcinoma cells. Glucosidase IIb was first purified from a human carcinoma cell line as a potential substrate for protein kinase C. Through the HDEL signal at the C-terminus, Glucosidase IIb retains the complete complex in the ER.

Function:

Cleaves sequentially the 2 innermost alpha-1,3-linked glucose residues from the Glc(2)Man(9)GlcNAc(2) oligosaccharide precursor of immature glycoproteins.

Subunit:

Heterodimer of a catalytic alpha subunit (GANAB) and a beta subunit (PRKCSH). Binds glycosylated PTPRC (By similarity).

Subcellular Location:

Endoplasmic reticulum. Golgi apparatus. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

Tissue Specificity:

Detected in placenta.

Similarity:

Belongs to the glycosyl hydrolase 31 family.

Database links:

[Entrez Gene: 23193](#) Human

[Omir: 104160](#) Human

[SwissProt: Q14697](#) Human

[Unigene: 595071](#) Human

Important Note:

This product as supplied is intended for research use only, not for use in



SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

human, therapeutic or diagnostic applications.