SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Marker Primary Antibody]

Rabbit Anti-SART1/AF647 Conjugated antibody

Catalog Number: SL7874R-AF647

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Target:	Anti-SART1/AF647
Immunogen:	KLH conjugated synthetic peptide derived from human SART1
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Applications:	IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	(predicted:Human,Mouse,Rat,Pig,Cow,Sheep)
Background:	background: SART1 encodes two proteins, the SART1(800) protein expressed in the nucleus of the majority of proliferating cells, and the SART1(259) protein expressed in the cytosol of epithelial cancers. The SART1(259) protein is translated by the mechanism of -1 frameshifting during posttranscriptional regulation; its full length sequence is not published yet. The two encoded proteins are thought to be involved in the regulation of proliferation. Both proteins have tumor rejection antigens. The SART1(259) protein possesses tumor epitopes capable of inducing HLA-A2402-restricted cytotoxic T lymphocytes in cancer patients. This SART1(259) antigen may be useful in specific immunotherapy for cancer patients and may serve as a paradigmatic tool for the diagnosis and treatment of patients with atopy. The SART1(259) protein is found to be essential for the recruitment of the tri snRNP to the pre spliceosome in the spliceosome assembly pathway. **Function:** May play a role in mRNA splicing. May also bind to DNA. **Subunit:** Identified in the spliceosome C complex. Part of a tri-snRNP

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

	complex. **Subcellular Location:** Nucleus. Note=Found in the nucleus of mitogen-activated peripheral blood mononuclear cells (PBMCs), tumor cells, or normal cell lines, but not in normal tissues except testis and fetal liver or in unstimulated PBMCs, suggesting preferential expression in proliferating cells. **Tissue Specificity:** Ubiquitously expressed. **Post-translational modifications:** Sumoylated with SUMO2. Phosphorylated upon DNA damage, probably by ATM or ATR. **Similarity:** Belongs to the SNU66/SART1 family. Important **Note:** This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.