

Rabbit Anti-hnRNP U/APC Conjugated antibody

SL9092R-APC

Product Name	Anti-hnRNP U/APC
Chinese Name	APC 标记的异质核糖核蛋白 U 抗体
Alias	Heterogeneous nuclear ribonucleoprotein U; hnRNP U; hnRNP U protein; HNRNPU; hnRNPU protein; HNRPU; HNRPU_HUMAN; p120; p120 nuclear protein; pp120; SAF A; SAF-A; SAFA; Scaffold attachment factor A; U21.1.
Research Area	Cell biology immunology Epigenetics
Immunogen Species	Rabbit
Clonality	Polyclonal
React Species	Human,Mouse(predicted:Rat,Pig,Cow)
Applications	IF=1:100-500, ICC/IF=1:50 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight	90kDa
Form	Lyophilized or Liquid
Concentration	1mg/ml
immunogen	KLH conjugated synthetic peptide derived from human hnRNP U
Lsotype	IgG
Purification	affinity purified by Protein A
Storage Buffer	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 1M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Product Detail	background: Heterogeneous nuclear ribonucleoproteins (hnRNPs) are thought to be involved in pre-mRNA processing. However, its role in the regulation of gene expression is as yet poorly understood. Proteins of the heterogeneous nuclear

ribonucleoparticles (hnRNP) family form a structurally diverse group of RNA binding proteins implicated in various functions. Recently, hnRNP proteins have been shown to hinder communication between factors bound to different splice sites. Conversely, several reports have described a positive role for some hnRNP proteins in pre-mRNA splicing. hnRNP-U, also termed scaffold attachment factor A (SAF-A), binds to pre-mRNA and nuclear matrix/scaffold attachment region DNA elements.

Function:

Component of the CRD-mediated complex that promotes MYC mRNA stabilization. Binds to pre-mRNA. Has high affinity for scaffold-attached region (SAR) DNA. Binds to double- and single-stranded DNA and RNA.

Subunit:

Identified in the spliceosome C complex. Component of the coding region determinant (CRD)-mediated complex, composed of DHX9, HNRNPU, IGF2BP1, SYNCRIP and YBX1. Identified in a mRNP complex, at least composed of DHX9, DDX3X, ELAVL1, HNRNPU, IGF2BP1, ILF3, PABPC1, PCBP2, PTBP2, STAU1, STAU2, SYNCRIP and YBX1. Identified in a mRNP granule complex, at least composed of ACTB, ACTN4, DHX9, ERG, HNRNPA1, HNRNPA2B1, HNRNPAB, HNRNPD, HNRNPL, HNRNPR, HNRNPU, HSPA1, HSPA8, IGF2BP1, ILF2, ILF3, NCBP1, NCL, PABPC1, PABPC4, PABPN1, RPLP0, RPS3, RPS3A, RPS4X, RPS8, RPS9, SYNCRIP, TROVE2, YBX1 and untranslated mRNAs. Interacts with IGF2BP1 and ERBB4. Ligand for CR2.

Subcellular Location:

Nucleus. Cytoplasm. Cell surface. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Component of ribonucleosomes. Also found associated with the cell surface.

Post-translational modifications:

Extensively phosphorylated.
Arg-733 and Arg-739 are dimethylated, probably to asymmetric dimethylarginine.

Similarity:

Contains 1 B30.2/SPRY domain.
Contains 1 SAP domain.

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com
