



Rabbit Anti-Proteasome 20S alpha 3(Ser250) antibody

SL9353R

Product Name:	Proteasome 20S alpha 3(Ser250)
Chinese Name:	磷酸化蛋白酶体PSM α 3抗体
Alias:	Proteasome 20S alpha 3 (phospho S250); 20S Proteasome α 3; HC8; Macropain subunit C8; Multicatalytic endopeptidase complex subunit C8; Proteasome (prosome macropain) subunit alpha type 3; Proteasome alpha 3 subunit; Proteasome component C8; Proteasome subunit alpha type 3; Proteasome subunit alpha type-3; Proteasome subunit C8; PSA3 HUMAN; PSC3; PSMA3.
Organism Species:	Rabbit
Clonality:	Polyclonal
React Species:	Human,Mouse,Rat,Dog,Pig,Cow,Horse,Rabbit,Sheep,
Applications:	ELISA=1:500-1000IHC-P=1:400-800IHC-F=1:400-800IF=1:50-200 (Paraffin sections need antigen repair) not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Molecular weight:	28kDa
Cellular localization:	The nucleuscytoplasmic
Form:	Lyophilized or Liquid
Concentration:	1mg/ml
immunogen:	KLH conjugated Synthesised phosphopeptide derived from human Proteasome 20S alpha 3 around the phosphorylation site of Ser250:DE(p-S)DD
Isotype:	IgG
Purification:	affinity purified by Protein A
Storage Buffer:	0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Storage:	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed:

[PubMed](#)

The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. Binds to the C-terminus of CDKN1A and thereby mediates its degradation. Negatively regulates the membrane trafficking of the cell-surface thromboxane A2 receptor (TBXA2R) isoform 2.

Function:

The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. Binds to the C-terminus of CDKN1A and thereby mediates its degradation. Negatively regulates the membrane trafficking of the cell-surface thromboxane A2 receptor (TBXA2R) isoform 2.

Subunit:

The 26S proteasome consists of a 20S proteasome core and two 19S regulatory subunits. The 20S proteasome core is composed of 28 subunits that are arranged in four stacked rings, resulting in a barrel-shaped structure. The two end rings are each formed by seven alpha subunits, and the two central rings are each formed by seven beta subunits. The catalytic chamber with the active sites is on the inside of the barrel. Interacts with AURKB. Interacts with CDKN1A. Interacts with HIV-1 TAT protein. Interacts with hepatitis C virus (HCV) F protein. Interacts with Epstein-Barr virus EBNA3 proteins. Interacts with MDM2 and RB1. Interacts with the C-terminus of TBXA2R isoform 2.

Subcellular Location:

Cytoplasm. Nucleus.

Similarity:

Belongs to the peptidase T1A family.

SWISS:

P25788

Gene ID:

5684

Database links:

[Entrez Gene: 5684](#)Human

[Entrez Gene: 19167](#)Mouse

[Entrez Gene: 29670](#)Rat

Product Detail:

[Entrez Gene: 408248](#)Rat

[Oimim: 176843](#)Human

[SwissProt: P25788](#)Human

[SwissProt: O70435](#)Mouse

[SwissProt: P18422](#)Rat

[Unigene: 558799](#)Human

[Unigene: 296338](#)Mouse

[Unigene: 3997](#)Rat

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

www.sunlongbiotech.com