SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Recombinant Rabbit Anti-PTPRC Monoclonal Antibody

Catalog Number: SLA0616M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Mouse
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	2A10
GeneID:	5788
SWISS:	P08575
Target:	PTPRC
Immunogen:	Recombinant human CD45 protein
Purification:	affinity purified by Protein A
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20℃ for one year. Avoid repeated freeze/thaw cycles.
Applications:	IHC-P (1:200-800) IHC-F (1:200-800) IF (1:200-800) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human
Predicted MW.:	143 kDa
Subcellular Location:	Membrane
Background:	The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus belongs to receptor type PTP. This gene is specifically expressed in hematopoietic cells. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SUNLONG

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

	receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling CD45 (LCA, leukocyte common antigen) is a receptor-type protein tyrosine phosphatase (PTP) ubiquitously expressed in all nucleated hematopoietic cells, comprising approximately 10% of all surface proteins in lymphocytes. CD45 is absent on non-hematopoietic cell lines, normal and malignant, non-hematopoietic tissues. CD45 glycoprotein is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases. CD45 protein exists as multiple isoforms as a result of alternative splicing, differ in their extracellular domains but share identical transmembrane and cytoplasmic domains. CD45RA is an isoform of the CD45 complex and has restricted expression between different subtypes of lymphoid cells. CD45 isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. CD45 has been shown to be an essential regulator of T- and B-cell antigen receptor signaling and suppresses JAK kinases to regulate cytokine receptor signaling. CD45 is also important in promoting cell survival by modulating integrin-mediated signal transduction pathway, DNA fragmentation during apoptosis and inhibition or upregulation of various immunological functions.

[VALIDATION IMAGES]




Figure 1. Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.




Figure 2. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com




Figure 3. Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.




Figure 4. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Lymph; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.




Figure 6. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.




Figure 7. Paraformaldehyde-fixed, paraffin embedded Human Small Intestine; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD45 Monoclonal Antibody, Unconjugated (SLA0616M) at 1:200 overnight at 4°C, followed by conjugation to the SL40296G-HRP and DAB (C-0010) staining.
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.