SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Recombinant Rabbit Anti-CDKN1B Monoclonal Antibody

Catalog Number: SLA0704M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Clone No.:	10B7
GeneID:	1027
SWISS:	P46527
Target:	CDKN1B
Immunogen:	Recombinant human CDKN1B
Purification:	affinity purified by Protein G
Concentration:	1mg/1ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20℃ for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human,Mouse,Rat
Predicted MW.:	22 kDa
Subcellular Location:	Nucleus
Background:	Cell cycle progression is regulated by cyclins and their cognate Cdks. p27 KIP 1 is a cell cycle regulatory mitotic inhibitor of cdk activity. p27 KIP 1 is a candidate tumor suppressor gene, and has been proposed to function as a possible mediator of TGF beta induced G1 arrest. p27 KIP 1 is up regulated in response to antimitogenic stimuli. The increased protein expression of p27 results in cellular arrest by binding to cyclin/Cdk complexes such as cyclin D1/Cdk4. p27 Kip1 is regulated by phosphorylation on serine 10 (S10) and threonine 187 (T187). Phosphorylation by CDK2 on T187 results in

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

	ubiquitylation and degradation of p27 Kip 1; while phosphorylation by hKIS on S10 signals the nuclear export to the cytoplasm.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.