SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Recombinant Rabbit Anti-CDKN2A Monoclonal Antibody

Catalog Number: SLA0773M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Mouse
Isotype:	IgG2b/Kappa
Clonality:	Monoclonal
Clone No.:	4E10
SWISS:	P42771
Target:	CDKN2A
Immunogen:	Synthesized peptide derived from human p16 INK
Purification:	affinity purified by Protein G
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human
Predicted MW.:	17 kDa
Background:	This gene generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported , two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with , and sequester , the E3 ubiquitin-protein ligase MDM2 , a protein responsible for the degradation of p53. In spite of the structural and

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

	functional differences , the CDK inhibitor isoforms and the ARF product encoded by this gene , through the regulatory roles of CDK4 and p53 in cell cycle G1 progression , share a common functionality in cell cycle G1 control. This gene is frequently mutated or deleted in a wide variety of tumors , and is known to be an important tumor suppressor gene. [provided by RefSeq , Sep 2012]

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.