SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

**[ Primary Antibody ]**

# Recombinant Rabbit Anti-CHI3L1 Monoclonal Antibody

**Catalog Number: SLA1048M**

**FOR RESEARCH USE ONLY**

**[ DATASHEET ]**

| | |
|---|---|
| **Host:** | Rabbit |
| **Isotype:** | IgG |
| **Clonality:** | Monoclonal |
| **Clone No.:** | 1H3 |
| **GeneID:** | 1116 |
| **SWISS:** | P36222 |
| **Target:** | CHI3L1 |
| **Immunogen:** | KLH conjugated synthetic peptide derived from human CHI3L1 |
| **Purification:** | affinity purified by Protein A |
| **Concentration:** | 1mg/ml |
| **Storage:** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20℃ for one year. Avoid repeated freeze/thaw cycles. |
| **Applications:** | WB (1:500-2000)<br>IHC-P (1:100-500)<br>IHC-F (1:100-500)<br>IF (1:100-500)<br>Not yet tested in other applications.<br>Optimal working dilutions must be determined by the end user. |
| **Reactivity:** | Human |
| **Predicted MW.:** | 42 kDa |
| **Background:** | Chitinases catalyze the hydrolysis of chitin, which is an abundant glycopolymer found in insect exoskeletons and fungal cell walls. The glycoside hydrolase 18 family of chitinases includes eight human family members. This gene encodes a glycoprotein member of the glycosyl hydrolase 18 family. The protein lacks chitinase activity and is secreted by activated macrophages, chondrocytes, neutrophils and synovial cells. The protein is thought to play a role in the process of inflammation and tissue remodeling. [provided by RefSeq, Sep 2009] |

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

## [ VALIDATION IMAGES ]





Figure 1. THP-1 (H) treated with PMA (100 ng/mL) for 24 h, Replace with fresh complete culture medium and continue to culture for 24 h, + hIFN-α (500U/mL) for 24 h, then with Brefeldin A (1μg/ml) for 3 h, 25 μg total protein per lane of cell lysates (see on figure) probed with CHI3L1 monoclonal antibody, unconjugated (SLA1048M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Figure 2. Paraformaldehyde-fixed, paraffin embedded Human Lung squamous cell carcinoma; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CHI3L1 Monoclonal Antibody, Unconjugated (SLA1048M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 3. Paraformaldehyde-fixed, paraffin embedded Human Glioma; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CHI3L1 Monoclonal Antibody, Unconjugated (SLA1048M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 4. Paraformaldehyde-fixed, paraffin embedded Human Ovarian Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CHI3L1 Monoclonal Antibody, Unconjugated (SLA1048M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CHI3L1 Monoclonal Antibody, Unconjugated (SLA1048M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 6. Paraformaldehyde-fixed, paraffin embedded Human Lung; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CHI3L1 Monoclonal Antibody, Unconjugated (SLA1048M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com




Figure 7. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CHI3L1 Monoclonal Antibody, Unconjugated (SLA1048M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.