SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

**[ Primary Antibody ]**

# Recombinant Rabbit Anti-BIRC2 Monoclonal Antibody

**Catalog Number: SLA1586M**

**FOR RESEARCH USE ONLY**

**[ DATASHEET ]**

| | |
|---|---|
| **Host:** | Rabbit |
| **Isotype:** | IgG |
| **Clonality:** | Monoclonal |
| **Clone No.:** | 16H6 |
| **GeneID:** | 329 |
| **SWISS:** | Q13490 |
| **Target:** | BIRC2 |
| **Immunogen:** | A synthesized peptide derived from human BIRC2 |
| **Purification:** | affinity purified by Protein A |
| **Storage:** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| **Applications:** | WB (1:500-2000)<br>IHC-P (1:100-500)<br>IHC-F (1:100-500)<br>IF (1:100-500)<br>Not yet tested in other applications.<br>Optimal working dilutions must be determined by the end user. |
| **Reactivity:** | Human |
| **Predicted MW.:** | 70 kDa |
| **Background:** | Multi-functional protein which regulates not only caspases and apoptosis, but also modulates inflammatory signaling and immunity, mitogenic kinase signaling, and cell proliferation, as well as cell invasion and metastasis. Acts as an E3 ubiquitin-protein ligase regulating NF-kappa-B signaling and regulates both canonical and non-canonical NF-kappa-B signaling by acting in opposite directions: acts as a positive regulator of the canonical pathway and suppresses constitutive activation of non-canonical NF-kappa-B signaling. |

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

## [ VALIDATION IMAGES ]





Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with cIAP1 monoclonal antibody, unconjugated (SLA1586M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Figure 2. Paraformaldehyde-fixed, paraffin embedded Human Lymph; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with cIAP1 Monoclonal Antibody, Unconjugated(SLA1586M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 3. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with cIAP1 Monoclonal Antibody, Unconjugated(SLA1586M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 4. Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with cIAP1 Monoclonal Antibody, Unconjugated(SLA1586M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Gastric Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with cIAP1 Monoclonal Antibody, Unconjugated(SLA1586M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.