SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Recombinant Rabbit Anti-CEACAM6 Monoclonal Antibody

Catalog Number: SLA1608M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	4E2
GeneID:	4680
SWISS:	P40199
Target:	CEACAM6
Immunogen:	A synthesized peptide derived from human CEACAM6
Purification:	affinity purified by Protein A
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:50-200) IP (1:20-50) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human
Predicted MW.:	37 kDa
Background:	Cell surface glycoprotein that plays a role in cell adhesion and tumor progression.Intercellular adhesion occurs in a calcium- and fibronectin-independent manner.Mediates homophilic and heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM5 and CEACAM8.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[VALIDATION IMAGES]




Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with CEACAM6 monoclonal antibody, unconjugated (SLA1608M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Figure 2. Paraformaldehyde-fixed, paraffin embedded Human lung;Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CEACAM6 Monoclonal Antibody, Unconjugated (SLA1608M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 3. Paraformaldehyde-fixed, paraffin embedded Human colon;Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CEACAM6 Monoclonal Antibody, Unconjugated (SLA1608M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 4. Paraformaldehyde-fixed, paraffin embedded Human thyroid tumor;Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CEACAM6 Monoclonal Antibody, Unconjugated (SLA1608M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human colon cancer;Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CEACAM6 Monoclonal Antibody, Unconjugated (SLA1608M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.