SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

**[ Primary Antibody ]**

# Recombinant Rabbit Anti-SMN1 Monoclonal Antibody

**Catalog Number: SLA1933M**

**FOR RESEARCH USE ONLY**

**[ DATASHEET ]**

| | |
|---|---|
| **Host:** | Rabbit |
| **Isotype:** | IgG |
| **Clonality:** | Monoclonal |
| **Clone No.:** | 7B6 |
| **GeneID:** | 6606 |
| **SWISS:** | Q16637 |
| **Target:** | SMN1 |
| **Immunogen:** | A synthesized peptide derived from human SMN1 |
| **Purification:** | affinity purified by Protein A |
| **Storage:** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| **Applications:** | WB (1:500-2000)<br>IHC-P (1:100-500)<br>IHC-F (1:100-500)<br>IF (1:100-500)<br>IP (1:20-50)<br>Not yet tested in other applications.<br>Optimal working dilutions must be determined by the end user. |
| **Reactivity:** | Human,Mouse,Rat |
| **Predicted MW.:** | 32 kDa |
| **Subcellular Location:** | Nucleus |
| **Background:** | The SMN complex catalyzes the assembly of small nuclear ribonucleoproteins (snRNPs), the building blocks of the spliceosome, and thereby plays an important role in the splicing of cellular pre-mRNAs.Most spliceosomal snRNPs contain a common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the core snRNP. |

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

## [ VALIDATION IMAGES ]





Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with SMN1 monoclonal antibody, unconjugated (SLA1933M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Figure 2. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMN1 Monoclonal Antibody, Unconjugated (SLA1933M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 3. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMN1 Monoclonal Antibody, Unconjugated (SLA1933M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 4. Paraformaldehyde-fixed, paraffin embedded Mouse Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMN1 Monoclonal Antibody, Unconjugated (SLA1933M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SMN1 Monoclonal Antibody, Unconjugated (SLA1933M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.