SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Recombinant Rabbit Anti-CHAF1B Monoclonal Antibody

Catalog Number: SLA1934M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	4H9
GeneID:	8208
SWISS:	Q13112
Target:	CHAF1B
Immunogen:	A synthesized peptide derived from human CAF 1 subunit B
Purification:	affinity purified by Protein A
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB (1:500-2000) IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200) Flow-Cyt (1:50-100) ICC/IF (1:50-200) IP (1:20-50) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human
Predicted MW.:	61 kDa
Subcellular Location:	N/A
Background:	Complex that is thought to mediate chromatin assembly in DNA replication and DNA repair. Assembles histone octamers onto replicating DNA in vitro. CAF-1 performs the first step of the nucleosome assembly process, bringing newly synthesized histones H3 and H4 to replicating DNA; histones H2A/H2B can bind to this chromatin precursor subsequent to DNA replication to complete the histone octamer.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

[VALIDATION IMAGES]




Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with p60 CAF1 monoclonal antibody, unconjugated (SLA1934M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min




Figure 2. Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with p60 CAF1 Monoclonal Antibody, Unconjugated (SLA1934M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 3. Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with p60 CAF1 Monoclonal Antibody, Unconjugated (SLA1934M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 4. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with p60 CAF1 Monoclonal Antibody, Unconjugated (SLA1934M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with p60 CAF1 Monoclonal Antibody, Unconjugated (SLA1934M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining




Figure 6. Paraformaldehyde-fixed, paraffin embedded Human Lymph; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with p60 CAF1 Monoclonal Antibody, Unconjugated (SLA1934M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com




Figure 7. The K562 (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C,the cells then were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.), followed by secondary antibody incubation for 40 min at room temperature. Primary Antibody (green):Rabbit Anti-p60 CAF1 antibody (SLA1934M,1:100); Isotype Control (orange): Rabbit IgG (SL0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed




Figure 8. 4% Paraformaldehyde-fixed K562 (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (p60 CAF1) monoclonal Antibody, unconjugated (SLA1934M) 1:100, 90 min at 37°C; followed by BF488 conjugated Goat Anti-Rabbit IgG antibody (green, SL60295G-BF488) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.