SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Recombinant Rabbit Anti-PIAS1&PIAS2 Monoclonal Antibody

Catalog Number: SLA2802M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Rabbit
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	10G6
GeneID:	8554
SWISS:	O75925
Target:	PIAS1&PIAS2
Immunogen:	A synthesized peptide derived from human PIAS1
Purification:	affinity purified by Protein A
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20℃ for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) IP (1:20-50) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human,Mouse,Rat
Predicted MW.:	72,68 kDa
Background:	Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53 pathway and the steroid hormone signaling pathway.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

[VALIDATION IMAGES]




Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with PIAS1/2 monoclonal antibody, unconjugated (SLA2802M) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Figure 2. Paraformaldehyde-fixed, paraffin embedded Rat Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 3. Paraformaldehyde-fixed, paraffin embedded Mouse Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 4. Paraformaldehyde-fixed, paraffin embedded Rat Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 5. Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 6. Paraformaldehyde-fixed, paraffin embedded Mouse Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com




Figure 7. Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 8. Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 9. Paraformaldehyde-fixed, paraffin embedded Rat Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 10. Paraformaldehyde-fixed, paraffin embedded Mouse Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 11. Paraformaldehyde-fixed, paraffin embedded Rat Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 12. Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com




Figure 13. Paraformaldehyde-fixed, paraffin embedded Mouse Spleen; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 14. Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Figure 15. Paraformaldehyde-fixed, paraffin embedded Rat testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Ratbbit IgG antibody (red, SL0295G-BF594), DAPI (blue, C02-04002) was used to stain the cell nuclei.




Figure 16. Paraformaldehyde-fixed, paraffin embedded Mouse testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with PIAS1/2 Monoclonal Antibody, Unconjugated (SLA2802M) at 1:200 overnight at 4°C. Followed by conjugated Goat Anti-Ratbbit IgG antibody (red, SL0295G-BF594), DAPI (blue, C02-04002) was used to stain the cell nuclei.
FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic procedures.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.