



PCNA Recombinant Rabbit mAb

SLM-33035R

Product Name PCNA Recombinant Rabbit mAb

Chinese Name 增殖 The nucleus 抗原（内参）Recombinant rabbit monoclonal anti

Alias Cyclin; DNA polymerase delta auxiliary protein; HGNC8729; MGC8367; Mutagen-sensitive 20
PCNAR; Polymerase delta accessory protein; Proliferating Cell Nuclear Antigen

Product Type Internal reference anti Recombinant rabbit monoclonal anti

**Immunogen
Species** Rabbit

Clonality Recombinant

Clone NO. 14D10

React Species Human,Mouse,Rat

WB=1:5000-20000,IHC-P=1:200-1000,IHC-F=1:200-1000,IF=1:200-1000

Applications not yet tested in other applications.
optimal dilutions/concentrations should be determined by the end user.

**Theoretical
molecular
weight** 29 kDa

**Detection
molecular
weight**

Form Liquid

Concentration 1mg/ml

immunogen KLH conjugated synthetic peptide derived from human PCNA

Lsotype IgG

Purification affinity purified by Protein A

**Buffer
Solution** 1M TBS (pH7.4) with 1% BSA, 2% Proclin300 and 50% Glycerol.


Storage Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Attention This product as supplied is intended for research use only, not for use in human, therapeutic or d


PubMed [PubMed](#)

**Product
Detail**


**Product
Picture**




25 ug total protein per lane of various lysates (see on figure) probed with PCNA monoclonal antibody (SLM-33035R) at 1:10000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.




Paraformaldehyde-fixed, paraffin embedded Human Spleen; Antigen retrieval by boiling in so (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Duodenum; Antigen retrieval by boiling (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated (C-0010) 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Colon Cancer; Antigen retrieval by boiling in citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated (C-0010) 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Liver Cancer; Antigen retrieval by boiling in citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated (C-0010) 1:200 overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Mouse Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Rat Kidney; Antigen retrieval by boiling in sodium citrate buffer for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SLM-3303) at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Mouse Kidney; Antigen retrieval by boiling in so (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in so (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Rat Colon; Antigen retrieval by boiling in sodium citrate buffer for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SLM-3303) at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Mouse Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Testicles; Antigen retrieval by boiling in (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Rat Testicles; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Testicles; Antigen retrieval by boiling in $\text{pH}6.0$ for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C , followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Rat Liver; Antigen retrieval by boiling in sodium citrate buffer for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SLM-3303) at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with PCNA Monoclonal Antibody, Unconjugated(SL0295G) overnight at 4°C, followed by conjugation to the SL0295G-HRP and DAB (C-0010) staining.