




## Mouse Anti-DNA-RNA Hybrid antibody

SLM-34118M

<b>Product Name</b>	DNA-RNA Hybrid
<b>Chinese Name</b>	DNA-RNA 杂交体单克隆抗体
<b>Alias</b>	DNA-RNA hybrid; RNA/DNA hybrid; DNA-RNA Hybrid [S9.6];
<b>Immunogen Species</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clone NO.</b>	S9.6
<b>React Species</b>	Species independent ICC/IF=1:50-200,ELISA=1:5000-10000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
<b>Applications</b>	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>immunogen</b>	ΦX174 bacteriophage-derived synthetic DNA/RNA
<b>Lsotype</b>	IgG2a
<b>Purification</b>	affinity purified by Protein A
<b>Buffer Solution</b>	PBS with 0.1% sodium azide and 50% glycerol, pH 7.4
<b>Storage</b>	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
<b>Attention</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>PubMed</b>	<a href="#">PubMed</a>
<b>Product Detail</b>	

**Product  
Picture**



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (DNA-RNA Hybrid) monoclonal Antibody, Unconjugated (SLM-34118M) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Mouse IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.