

## Mouse Anti-Procalcitonin (PCT)antibody

SLM-41284M

<b>Product Name</b>	Procalcitonin (PCT)
<b>Chinese Name</b>	降钙素原单克隆抗体
<b>Alias</b>	Calcitonin; CALCA; Calcitonin carboxyl-terminal peptide; CCP; Katalcalcin; PDN-21; CALC1;
<b>Research Area</b>	Cardiovascular immunology Signal transduction Growth factors and hormones
<b>Immunogen Species</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>React Species</b>	Human
<b>Applications</b>	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500 not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Theoretical molecular weight</b>	13kDa
<b>Cellular localization</b>	Secretory protein
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>immunogen</b>	KLH conjugated synthetic peptide derived from human Procalcitonin(PCT)
<b>Lsotype</b>	IgG
<b>Purification</b>	affinity purified by Protein A
<b>Buffer Solution</b>	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
<b>Storage</b>	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
<b>Attention</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>PubMed</b>	<a href="#">PubMed</a>
<b>Product Detail</b>	Procalcitonin (PCT) is a peptide precursor of the hormone calcitonin, the latter being involved with calcium homeostasis. It arises once preprocalcitonin is cleaved by endopeptidase. It is composed of 116 amino acids and is produced

by parafollicular cells (C cells) of the thyroid and by the neuroendocrine cells of the lung and the intestine.

**Function:**

The level of procalcitonin in the blood stream of healthy individuals is below the limit of detection (1 µg/L) of clinical assays. The level of procalcitonin rises in a response to a proinflammatory stimulus, especially of bacterial origin. It does not rise significantly with viral or non-infectious inflammations. With the derangements that a severe infection with an associated systemic response brings, the blood levels of procalcitonin may rise to 100 µg/L. In serum, procalcitonin has a half-life of 25 to 30 hours. Remarkably the high procalcitonin levels produced during infections are not followed by a parallel increase in calcitonin or a decrease in serum calcium levels.

**DISEASE:**

Sepsis: Measurement of procalcitonin can be used as a marker of severe sepsis caused by bacteria and generally grades well with the degree of sepsis, although levels of procalcitonin in the blood are very low. PCT has the greatest sensitivity (85%) and specificity (91%) for differentiating patients with systemic inflammatory response syndrome (SIRS) from those with sepsis, when compared with IL-2, IL-6, IL-8, CRP and TNF-alpha. Evidence is emerging that procalcitonin levels can reduce unnecessary antibiotic prescribing to people with lower respiratory tract infections. Currently, procalcitonin assays are widely used in the clinical environment.

Pneumonia: Procalcitonin levels may be useful to distinguish bacterial infections from nonbacterial infections. This may help guide antibiotic use, which can help save on cost and drug resistance.

Kidney disease: Patients with chronic kidney disease and end-stage renal disease are at higher risk for infections, and procalcitonin has been studied in these populations, who often have higher levels. Procalcitonin can be dialyzed, and so levels are dependent upon when patients receive hemodialysis. While there is no formally accepted cutoff value for patients undergoing HD, using a value of greater or equal to 0.5 ng/mL yielded a sensitivity of 97-98% and a specificity of 70-96%.

Hepatitis: PCT, possibly together with CRP, is used to corroborate the MELD score.

**Similarity:**

Belongs to the calcitonin family.

**SWISS:**

P01258

**Gene ID:**

796

**Database links:**


[Entrez Gene: 796](#) Human

[Omim: 114130](#) Human

[SwissProt: P01258](#) Human

[Unigene: 37058](#) Human

**Product Picture**



Sample:

Lane 1: Recombinant human Procalcitonin, N-His

Primary: Anti-Procalcitonin (PCT) (SLM-41284M) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution



Predicted band size: 13 kDa

Observed band size: 17 kDa