

## Mouse Anti-MMP3 antibody

SLM-43017M

<b>Product Name</b>	MMP3
<b>Chinese Name</b>	基质金属蛋白酶 3 单克隆抗体
<b>Alias</b>	MMP3_HUMAN; Stromelysin-1; EC:3.4.24.17; STMY1; SL-1; SL 1; SL1; Matrix metalloproteinase-3 (MMP-3); MMP-3; MMP 3; Transin-1;
<b>Research Area</b>	Tumour
<b>Immunogen Species</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>React Species</b>	Human, WB=1:500-2000
<b>Applications</b>	not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
<b>Theoretical molecular weight</b>	52kDa
<b>Cellular localization</b>	Extracellular matrix Secretory protein
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>immunogen</b>	Recombinant human MMP-3 protein: 18-477/477
<b>Lsotype</b>	IgG
<b>Purification</b>	affinity purified by Protein A
<b>Buffer Solution</b>	1M TBS(pH7.4) with 1% BSA, 3% Proclin300 and 50% Glycerol.
<b>Storage</b>	Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
<b>Attention</b>	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
<b>PubMed</b>	<a href="#">PubMed</a>
<b>Product Detail</b>	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as

inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq, Jul 2008].

**Function:**

Can degrade fibronectin, laminin, gelatins of type I, III, IV, and V; collagens III, IV, X, and IX, and cartilage proteoglycans. Activates procollagenase.

**Subcellular Location:**

Secreted, extracellular space, extracellular matrix (Probable).

**Similarity:**

Belongs to the peptidase M10A family. Contains 4 hemopexin-like domains.

**SWISS:**

P08254

**Gene ID:**

4314

**Database links:**

[Entrez Gene: 4314](#) Human

[Entrez Gene: 17392](#) Mouse

[Entrez Gene: 171045](#) Rat

[Omim: 185250](#) Human

[SwissProt: P08254](#) Human

[SwissProt: P28862](#) Mouse


[SwissProt: P03957](#) Rat

[Unigene: 375129](#) Human

Synthesis and Degradation (Synthesis and Degradation) 基质金属蛋白酶 (matrix metalloproteinases, MMPs)是一族依赖锌离子而降解各种 Extracellular matrix 的蛋白酶, 亦称 IV 型胶原酶或称明胶酶 A, 其主要功能为降解 IV 型胶原, 因而它在 Tumour 细胞突破基底膜屏障和浸润转

移中起重要作用。MMP3 目前主要用于各种恶性 Tumour(如乳腺癌、胃肠道癌、卵巢癌、膀胱癌等)中的基底膜检测与 Tumour 转移浸润的研究。

**Product Picture**



Sample:

Lane 1: Recombinant human MMP-3 protein

Primary: Anti-MMP3 (SLM-43017M) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution

Predicted band size: 52 kDa

Observed band size: 60 kDa