SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Mouse Anti-Symmetric Di-Methyl-Histone H3 (Arg2)antibody

Catalog Number: SLM-60158M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Mouse
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	G3G8
Target:	Symmetric Di-Methyl-Histone H3 (Arg2)
Purification:	Antigen affinity purification
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB (1:500-2000) Not yet tested in other applications. Optimal working dilutions must be determined by the end user.
Reactivity:	Human, Rat (predicted: Mouse, Monkey)
Subcellular Location:	The nucleus
Background:	Histone post-translational modifications (PTMs) are a key mechanism by which epigenetics regulates chromatin structure and are known as the "histone code". Post-translational modifications on histones include acetylation, methylation, phosphorylation and some novel acylation modifications discovered in recent years. These histone modifications directly affect the binding of chromatin to transcription factors or other epigenetic regulators, and change genome stability and gene transcription, etc. Histone methylation usually occurs on lysine and arginine residues of core histones. Histone methylation can either promote or inhibit gene transcription, depending on whether histone methylation occurs on lysine or arginine and the number of methyl groups (lysine can undergo mono-, di- and tri-methylation, and arginine can undergo mono-, symmetric and asymmetric di-methylation). Histone lysine methylation usually occurs on lysine 4, 9, 27, 36 and 79 of histone H3 and lysine 20 of histone H4; arginine methylation usually occurs on arginine 2, 8, 17 and 26 of histone H3 and arginine 3 of histone H4. Protein methyltransferases (HMT) and histone demethylases (HDM) are the

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

SunLong Biotech Co.,LTD

Tel: 0086-571-56623320 Fax:0086-571-56623318

E-mail:sales@sunlongbiotech.com

www.sunlongbiotech.com

	main regulatory factors of methylation modifications.

[VALIDATION IMAGES]




Figure 1. 25 ug total protein per lane of various lysates (see on figure) probed with Symmetric Di-Methyl-Histone H3 (Arg2) monoclonal antibody, unconjugated (SLM-60158M) at 1:500 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.