SunLong Biotech Co.,LTD
Tel: 0086-571-56623320 Fax:0086-571-56623318
E-mail:sales@sunlongbiotech.com
www.sunlongbiotech.com

[Primary Antibody]

Mouse Anti-Butyryl-Histone H3 (Lys23)antibody

Catalog Number: SLM-60172M

FOR RESEARCH USE ONLY

[DATASHEET]

Host:	Mouse
Isotype:	IgG
Clonality:	Monoclonal
Clone No.:	H2B12
Target:	Butyryl-Histone H3 (Lys23)
Purification:	Antigen affinity purification
Concentration:	1mg/ml
Storage:	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4℃. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.
Applications:	WB=1:500-2000 IHC-P=1:100-500 IHC-F=1:100-500 IF=1:100-500（Paraffin sections need antigen repair） not yet tested in other applications. optimal dilutions/concentrations should be determined by the end user.
Reactivity:	Human(predicted:Mouse,Rat)
Subcellular Location:	The nucleus
Background:	Histones undergo various enzyme-catalyzed modifications, including acetylation, methylation, phosphorylation, ubiquitination, etc. Lysine butyrylation is a newly discovered reversible modification that controls chromosome structure and gene transcription. Lysine butyrylation is highly conserved in eukaryotic cells from worms to humans. The unique structural and genomic localization of histone lysine butyrylation suggests that it is mechanistically and functionally distinct from histone lysine acetylation. Specifically, in the genomes of human somatic cells and rabbit male germ cells, histone butyrylation marks active promoters or potential enhancers. Butyrylation of histone H3 at Lys18 may play an important role in epigenetic regulation, including chromatin remodeling and DNA transcriptional regulation.

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.